Colloquia: IFAE 2011

Neutrino physics: Status and perspectives

C. Giunti

INFN, Sezione di Torino - Via P. Giuria 1, I-10125 Torino, Italy

(ricevuto il 29 Luglio 2011; pubblicato online il 2 Novembre 2011)

Summary. — After a brief review of the results of solar, atmospheric and longbaseline neutrino oscillation experiments, which led to the current three-neutrino mixing paradigm, we discuss indications of neutrino oscillation experiments in favor of short-baseline oscillations which require the existence of one or more sterile neutrinos. We show that the simplest possibility of existence of one sterile neutrino is not enough to fit all data of short-baseline neutrino oscillation experiments because of two tensions: a tension between neutrino and antineutrino data and a tension between appearance and disappearance data.

PACS 14.60.Pq – Neutrino mass and mixing. PACS 14.60.Lm – Ordinary neutrinos. PACS 14.60.St – Non-standard-model neutrinos, right-handed neutrinos, etc.

The results of several solar, atmospheric and long-baseline neutrino oscillation experiments have proved that neutrinos are massive and mixed particles (see ref. [1]). There are two groups of experiments which measured two independent squared-mass differences (Δm^2) in two different neutrino flavor transition channels.

Solar neutrino experiments (Homestake, Kamiokande, GALLEX/GNO, SAGE, Super-Kamiokande, SNO, BOREXino) measured $\nu_e \rightarrow \nu_{\mu}, \nu_{\tau}$ oscillations generated by $\Delta m_{\rm SOL}^2 = 6.2^{+1.1}_{-1.9} \times 10^{-5} \, {\rm eV}^2$ and a mixing angle $\tan^2 \vartheta_{\rm SOL} = 0.42^{+0.04}_{-0.02}$ [2]. The KamLAND experiment confirmed these oscillations by observing the disappearance of reactor $\bar{\nu}_e$ at an average distance of about 180 km. The combined fit of solar and KamLAND data leads to $\Delta m_{\rm SOL}^2 = (7.6 \pm 0.2) \times 10^{-5} \, {\rm eV}^2$ and a mixing angle $\tan^2 \vartheta_{\rm SOL} = 0.44 \pm 0.03$ [2]. Notice that the agreement of solar and KamLAND data in favor of ν_e and $\bar{\nu}_e$ disappearance generated by the same oscillation parameters is consistent with the equality of neutrino and antineutrino disappearance expected from CPT symmetry (see ref. [1]).

Atmospheric neutrino experiments (Kamiokande, IMB, Super-Kamiokande, MACRO, Soudan-2, MINOS) measured ν_{μ} and $\bar{\nu}_{\mu}$ disappearance through oscillations generated by $\Delta m_{\rm ATM}^2 \simeq 2.3 \times 10^{-3} \, {\rm eV}^2$ and a mixing angle $\sin^2 2\vartheta_{\rm ATM} \simeq 1$ [3]. The K2K and MINOS long-baseline experiments confirmed these oscillations by observing the disappearance of accelerator ν_{μ} at distances of about 250 km and 730 km, respectively. The MINOS data give $\Delta m_{\rm ATM}^2 = 2.32^{+0.12}_{-0.08} \times 10^{-3} \, {\rm eV}^2$ and $\sin^2 2\vartheta_{\rm ATM} > 0.90$ at 90% CL [4]. The equality

© Società Italiana di Fisica

of muon neutrino and antineutrino disappearance expected from CPT symmetry is currently under investigation in the MINOS experiment [5], with preliminary results which hint at an intriguing difference between the muon neutrino and antineutrino oscillation parameters.

These measurements led to the current three-neutrino mixing paradigm, in which the three active neutrinos ν_e , ν_{μ} , ν_{τ} are superpositions of three massive neutrinos ν_1 , ν_2 , ν_3 with respective masses m_1 , m_2 , m_3 . The two measured squared-mass differences can be interpreted as $\Delta m_{\rm SOL}^2 = \Delta m_{21}^2$ and $\Delta m_{\rm ATM}^2 = |\Delta m_{31}^2| \simeq |\Delta m_{32}^2|$, with $\Delta m_{kj}^2 = m_k^2 - m_j^2$. In the standard parameterization of the 3 × 3 unitary mixing matrix (see ref. [1]) $\vartheta_{\rm SOL} \simeq \vartheta_{12}$, $\vartheta_{\rm ATM} \simeq \vartheta_{23}$ and $\sin^2 \vartheta_{13} < 0.035$ at 90% CL [6].

The completeness of the three-neutrino mixing paradigm has been challenged by the recent observation of a signal of short-baseline $\bar{\nu}_{\mu} \rightarrow \bar{\nu}_{e}$ oscillations in the MiniBooNE experiment [7] which agrees with a similar signal observed several years ago in the LSND experiment [8]. It is remarkable that the two signals have been observed at different values of distance (L) and energy (E), but approximately at the same L/E. Since the distance and energy dependences of neutrino oscillations occur through this ratio, the agreement of the MiniBooNE and LSND signals raised interest in the possibility of existence of one or more squared-mass differences much larger than Δm_{SOL}^2 and Δm_{ATM}^2 . These new squared-mass differences should have values larger than about 0.5 eV.

In the following, I consider the simplest extension of three-neutrino mixing with the addition of one massive neutrino. In such four-neutrino mixing framework the flavor neutrino basis is composed by the three active neutrinos ν_e , ν_{μ} , ν_{τ} and a sterile neutrino ν_s which does not have weak interactions.

The so-called 2+2 four-neutrino mixing schemes are strongly disfavored by the absence of any signal of sterile neutrino effects in solar and atmospheric neutrino data [9]. Hence, we must consider the so-called 3+1 four-neutrino schemes, in which the effective flavor transition and survival probabilities in short-baseline (SBL) experiments are given by

(1)
$$P^{\text{SBL}}_{\substack{(-)\\\nu_{\alpha}\to\nu_{\beta}}} = \sin^2 2\vartheta_{\alpha\beta}\sin^2\left(\frac{\Delta m^2 L}{4E}\right) \qquad (\alpha \neq \beta),$$

(2)
$$P_{\substack{(-)\\\nu_{\alpha}\to\nu_{\alpha}}}^{\text{SBL}} = 1 - \sin^2 2\vartheta_{\alpha\alpha} \sin^2 \left(\frac{\Delta m^2 L}{4E}\right),$$

for $\alpha, \beta = e, \mu, \tau, s$, with $\Delta m^2 = \Delta m_{\rm SBL}^2$ and

(3)
$$\sin^2 2\vartheta_{\alpha\beta} = 4|U_{\alpha4}|^2|U_{\beta4}|^2, \quad \sin^2 2\vartheta_{\alpha\alpha} = 4|U_{\alpha4}|^2\left(1-|U_{\alpha4}|^2\right).$$

Therefore:

- 1. All effective SBL oscillation probabilities depend only on the largest squared-mass difference $\Delta m^2 = \Delta m^2_{\text{SBL}} = |\Delta m^2_{41}|$.
- 2. All oscillation channels are open, each one with its own oscillation amplitude.
- 3. All oscillation amplitudes depend only on the absolute values of the elements in the fourth column of the mixing matrix, *i.e.* on three real numbers with sum less than unity, since the unitarity of the mixing matrix implies $\sum_{\alpha} |U_{\alpha 4}|^2 = 1$.



Fig. 1. – Exclusion curves obtained from the data of reactor $\bar{\nu}_e$ disappearance experiments (see ref. [10]), from the data of the CDHSW ν_{μ} disappearance experiment [11], and from atmospheric neutrino data (extracted from the analysis in ref. [12]).

4. *CP* violation cannot be observed in SBL oscillation experiments, even if the mixing matrix contains *CP*-violating phases. In other words, neutrinos and antineutrinos have the same effective SBL oscillation probabilities.

Before the recent indication of an antineutrino $\bar{\nu}_{\mu} \rightarrow \bar{\nu}_{e}$ signal consistent with the LSND antineutrino signal, the MiniBooNE Collaboration published the results of neutrino data which do not show a corresponding $\nu_{\mu} \rightarrow \nu_{e}$ signal [13]. This difference between the MiniBooNE neutrino and antineutrino data may be due to *CP* violation.

The absence of any difference in the effective SBL oscillation probabilities of neutrinos and antineutrinos in 3+1 four-neutrino mixing schemes implies that these schemes cannot explain the difference between neutrinos and antineutrino oscillations observed in the MiniBooNE. Moreover, the dependence of all the oscillation amplitudes in eq. (3) on three independent absolute values of the elements in the fourth column of the mixing matrix implies that the amplitude of $\stackrel{(-)}{\nu_{\mu}} \rightarrow \stackrel{(-)}{\nu_{e}}$ transitions is limited by the absence of large SBL disappearance of $\stackrel{(-)}{\nu_{e}}$ and $\stackrel{(-)}{\nu_{\mu}}$ observed in several experiments.

The results of reactor neutrino experiments constrain the value $|U_{e4}|^2$ through the measurement of $\sin^2 2\vartheta_{ee}$. The calculation of the reactor $\bar{\nu}_e$ flux has been recently improved in ref. [14], resulting in an increase of about 3% with respect to the previous value adopted by all experiments for the comparison with the data. The measured reactor rates are in agreement with those derived from the old $\bar{\nu}_e$ flux, but show a deficit of about 2.2 σ with respect to the rates derived from the new $\bar{\nu}_e$ flux. This is the "reactor antineutrino anomaly" [10](¹), which may be an indication in the $\bar{\nu}_e \rightarrow \bar{\nu}_e$ channel of a signal corresponding to the $\bar{\nu}_{\mu} \rightarrow \bar{\nu}_e$ signal observed in the LSND and MiniBooNE experiments. However, the $\bar{\nu}_e$ disappearance is small and large values of $|U_{e4}|^2$ close

 $[\]binom{1}{1}$ We do not consider here the "Gallium neutrino anomaly" [15-20], which may be compatible with the reactor antineutrino anomaly assuming the equality of neutrino and antineutrino disappearance imposed by the CPT symmetry.



Fig. 2. – (Colour on-line) Left panel: exclusion curves in the $\sin^2 2\vartheta_{e\mu}$ - Δm^2 plane obtained from the separate constraints in fig. 1 (blue and green lines) and the combined constraint given by eq. (6) (red line) from disappearance experiments (Dis). Right panel: exclusion curve obtained with the addition of KARMEN [21] (KAR), NOMAD [22] (NOM) and MiniBooNE neutrino [13] (MB ν) data (red line). In both panels the region enclosed by the dark-red lines is allowed by LSND and MiniBooNE antineutrino data.

to unity are excluded by solar neutrino oscillations (which require large $|U_{e1}|^2 + |U_{e2}|^2$), for small $\sin^2 2\vartheta_{ee}$ we have

(4)
$$\sin^2 2\vartheta_{ee} \simeq 4|U_{e4}|^2.$$

The value of $\sin^2 2\vartheta_{\mu\mu}$ is constrained by the curves in the right panel of fig. 1, which have been obtained from the lack of ν_{μ} disappearance in the CDHSW ν_{μ} experiment [11] and from the requirement of large $|U_{\mu1}|^2 + |U_{\mu2}|^2 + |U_{\mu3}|^2$ for atmospheric neutrino oscillations [12]. Hence, $|U_{\mu4}|^2$ is small and

(5)
$$\sin^2 2\vartheta_{\mu\mu} \simeq 4|U_{\mu4}|^2.$$

From eqs. (3), (4) and (5), for the amplitude of $\stackrel{(-)}{\nu_{\mu}} \rightarrow \stackrel{(-)}{\nu_{e}}$ transitions we obtain

(6)
$$\sin^2 2\vartheta_{e\mu} \simeq 0.25 \sin^2 2\vartheta_{ee} \sin^2 2\vartheta_{\mu\mu}.$$

Therefore, if $\sin^2 2\vartheta_{ee}$ and $\sin^2 2\vartheta_{\mu\mu}$ are small, $\sin^2 2\vartheta_{e\mu}$ is quadratically suppressed. This is illustrated in the left panel of fig. 2, where one can see that the separate effects of the constraints on $\sin^2 2\vartheta_{ee}$ and $\sin^2 2\vartheta_{\mu\mu}$ exclude only the large- $\sin^2 2\vartheta_{e\mu}$ part of the region allowed by LSND and MiniBooNE antineutrino data, whereas most of this region is excluded by the combined constraint in eq. (6). As shown in the right panel of fig. 2, the constraint becomes stronger by including the data of the KARMEN [21], NOMAD [22] and MiniBooNE neutrino [13] experiments, which did not observe a shortbaseline $\overset{(-)}{\nu_{\mu}} \rightarrow \overset{(-)}{\nu_{e}}$ signal. Since the parameter goodness of fit [23] is 0.0016%, 3+1 schemes are strongly disfavored by the data. This conclusion has been reached recently also in refs. [12, 24-26] and confirms the pre-MiniBooNE results in refs. [9, 27]. The CP-violating difference between MiniBooNE neutrino and antineutrino data can be explained by introducing another physical effect in addition to a sterile neutrino: a second sterile neutrino in 3+2 schemes [12, 24, 26, 28-31], non-standard interactions [24], CPT violation [25,32]. These additional effects also help in reducing the tension between appearance and disappearance data.

In conclusion, I think that we are living an exciting time in neutrino physics which may prelude to a transition from the well-established three-neutrino mixing paradigm to a new paradigm of neutrino mixing with sterile neutrinos and possibly other effects (as non-standard interactions and CPT violation) which are very interesting for the exploration of the physics beyond the Standard Model. In order to clarify the validity of the experimental indications in favor of an expansion of neutrino mixing beyond the standard three-neutrino mixing and resolve the tension between the current positive and negative experimental results, new experiments with high sensitivity and low background are needed.

REFERENCES

- [1] GIUNTI C. and KIM C. W., Fundamentals of Neutrino Physics and Astrophysics (Oxford University Press, Oxford, UK) 2007.
- [2] ABE K. et al. (SUPER-KAMIOKANDE), Phys. Rev. D, 83 (2011) 052010, arXiv:1010.0118.
- [3] ASHIE Y. et al. (SUPER-KAMIOKANDE), Phys. Rev. D, 71 (2005) 112005, hep-ex/0501064.
- [4] ADAMSON P. et al. (MINOS), Phys. Rev. Lett., **106** (2011) 181801, arXiv:1103.0340.
- [5] ADAMSON P. et al. (MINOS), arXiv:1104.0344 (2011).
- [6] SCHWETZ T., TORTOLA M. and VALLE J. W. F., New J. Phys., 10 (2008) 113011, arXiv:0808.2016.
- [7] AGUILAR-AREVALO A. A. et al. (MINIBOONE), Phys. Rev. Lett., 105 (2010) 181801, arXiv:1007.1150.
- [8] AGUILAR A. et al. (LSND), Phys. Rev. D, 64 (2001) 112007, hep-ex/0104049.
- [9] MALTONI M. et al., New J. Phys., 6 (2004) 122, hep-ph/0405172.
- [10] MENTION G. et al., Phys. Rev. D, 83 (2011) 073006, arXiv:1101.2755.
- [11] DYDAK F. et al. (CDHSW), Phys. Lett. B, 134 (1984) 281.
- [12] MALTONI M. and SCHWETZ T., Phys. Rev. D, 76 (2007) 093005, arXiv:0705.0107.
- [13] AGUILAR-AREVALO A. A. (MINIBOONE), Phys. Rev. Lett., 102 (2009) 101802, arXiv: 0812.2243.
- [14] MUELLER T. A. et al., Phys. Rev. C, 83 (2011) 054615, arXiv:1101.2663.
- [15] GIUNTI C. and LAVEDER M., Mod. Phys. Lett. A, 22 (2007) 2499, hep-ph/0610352.
- [16] GIUNTI C. and LAVEDER M., Phys. Rev. D, 77 (2008) 093002, arXiv:0707.4593.
- [17] ACERO M. A., GIUNTI C. and LAVEDER M., Phys. Rev. D, 78 (2008) 073009, arXiv: 0711.4222.
- [18] GIUNTI C. and LAVEDER M., Phys. Rev. D, 80 (2009) 013005, arXiv:0902.1992.
- [19] GIUNTI C. and LAVEDER M., Phys. Rev. D, 82 (2010) 053005, arXiv:1005.4599.
- [20] GIUNTI C. and LAVEDER M., Phys. Rev. C, 83 (2011) 065504, arXiv:1006.3244.
- [21] ARMBRUSTER B. et al. (KARMEN), Phys. Rev. D, 65 (2002) 112001, hep-ex/0203021.
- [22] ASTIER P. et al. (NOMAD), Phys. Lett. B, 570 (2003) 19, hep-ex/0306037.
- [23] MALTONI M. and SCHWETZ T., Phys. Rev. D, 68 (2003) 033020, hep-ph/0304176.
- [24] AKHMEDOV E. and SCHWETZ T., JHEP, 10 (2010) 115, arXiv:1007.4171.
- [25] GIUNTI C. and LAVEDER M., Phys. Rev. D, 83 (2011) 053006, arXiv:1012.0267.
- [26] KOPP J., MALTONI M. and SCHWETZ T., (2011) arXiv:1103.4570.
- [27] MALTONI M. et al., Nucl. Phys. B, 643 (2002) 321, hep-ph/0207157.
- [28] SOREL M., CONRAD J. and SHAEVITZ M., Phys. Rev. D, 70 (2004) 073004, hep-ph/ 0305255.
- [29] KARAGIORGI G. et al., Phys. Rev. D, 75 (2007) 013011, hep-ph/0609177.

- [30] KARAGIORGI G. et al., Phys. Rev. D, 80 (2009) 073001, arXiv:0906.1997.
 [31] GIUNTI C., talks presented at La Thuile 2011, NeuTel 2011 and IFAE 2011 (2011), arXiv:1106.4479.
- [32] GIUNTI C. and LAVEDER M., Phys. Rev. D, 82 (2010) 093016, arXiv:1010.1395.