

Top differential cross section measurements (Tevatron)

A. W. JUNG for the DØ COLLABORATION
Fermi National Accelerator Laboratory (Fermilab) - USA

ricevuto l' 1 Marzo 2012
pubblicato online il 28 Maggio 2012

Summary. — Differential cross sections in the top quark sector measured at the Fermilab Tevatron collider are presented. CDF used 2.7 fb^{-1} of data and measured the differential cross section as a function of the invariant mass of the $t\bar{t}$ system. The measurement shows good agreement with the standard model and furthermore is used to derive limits on the ratio κ/M_{Pl} for gravitons which decay to top quarks in the Randall-Sundrum model. DØ used 1.0 fb^{-1} of data to measure the differential cross section as a function of the transverse momentum of the top quark. The measurement shows a good agreement to the higher order perturbative QCD prediction and various predictions based on various Monte Carlo generators.

PACS 14.65.Ha – Top quarks.

1. – Introduction

The *top* quark is the heaviest known elementary particle and was discovered at the Tevatron $p\bar{p}$ collider in 1995 by the CDF and DØ collaboration [1, 2] at a mass of around 175 GeV. The production is dominated by the $q\bar{q}$ annihilation process with 85% as opposed to gluon-gluon fusion which contributes only 15%. Both measurements presented here are performed using the l +jets channel, where one of the W bosons (stemming from the decay of the *top* quarks) decays leptonically. The other W boson decays hadronically. The l +jets channel is a good compromise between signal and background contribution whilst having high event statistics. The branching fraction for top quarks decaying into Wb is almost 100%. Jets containing a beauty quark are identified by means of a neural network (NN) built by the combination of variables describing the properties of secondary vertices and of tracks with large impact parameters relative to the primary vertex.

2. – Measurement of the transverse momentum distribution of the top quarks

The measurement of the transverse momentum distribution of the top quarks [3] selects events with an isolated lepton with a transverse momentum p_T of at least 20 GeV and a pseudo-rapidity of $|\eta| < 1.1$ (e +jets) or $|\eta| < 2.0$ (μ +jets). A cut on the missing




Fig. 1. – Panel a) compares the background-subtracted reconstructed top-quark p_T distribution [3] with the one corrected for the effects of finite experimental resolution (two entries per event). Inner error bars represent the statistical uncertainty, whereas the outer one is statistical and systematic added in quadrature. Panel b) shows the unfolded invariant mass distribution of the $t\bar{t}$ system [4] compared to signal $t\bar{t}$ MC.

transverse energy (\cancel{E}_T) of 20 GeV is applied. Furthermore at least four jets are required with $p_T > 20$ GeV and $|\eta| < 2.5$, an additional cut of $p_T > 40$ GeV is applied for the leading jet. Finally at least one jet needs to be identified as a b -jet. For the reconstruction of the event kinematics additional constraints are used: the masses of the two W bosons are constrained to 80.4 GeV. Furthermore the masses of the two reconstructed top quarks are assumed to be equal. All possible permutations of objects are considered where the final solution is the one with the smallest χ^2 .

Figure 1a) shows the background-subtracted reconstructed top-quark p_T distribution compared to the one corrected for finite experimental resolution. The latter is derived by using regularized matrix unfolding. Figure 1b) compares the unfolded invariant mass distribution of the $t\bar{t}$ system to the expectation using $t\bar{t}$ signal MC. The correction for finite detector resolution is again done using regularized unfolding.

Figure 2a) shows the differential cross section as a function of top-quark p_T , where the leptonic and hadronic decay of the W boson to the top-quark cross section are combined. All predictions use the proton parton density function (PDF) CTEQ61 with the scale set to $\mu_r = \mu_f = m_t$ ($m_t = 170$ GeV) except for the approximate NNLO perturbative QCD (pQCD) prediction which uses the MSTW08 PDF. The normalization is nicely described by pQCD in (N)NLO, however there is an offset for PYTHIA and ALPGEN in normalization. Figure 2b) shows that the shape is reasonable described by all predictions. The inclusive total cross section for $t\bar{t}$ production is measured to $\sigma = 8.31 \pm 1.28(\text{stat.})$ pb and in good agreement with the latest theoretical predictions of $\sigma = 6.41 \pm_{0.42}^{0.51}$ pb [5] and $\sigma = 7.46 \pm_{0.67}^{0.48}$ pb [6].

3. – Measurement of the invariant mass distribution of the $t\bar{t}$ system

This measurement of the invariant mass distribution of the $t\bar{t}$ system $M_{t\bar{t}}$ [4] selects events with an isolated lepton with a p_T of at least 20 GeV and a pseudo-rapidity of




Fig. 2. – a) Differential cross section data (points) as a function of top-quark p_T (two entries per event) [3] compared with expectations from NLO pQCD (solid lines), from an approximate NNLO pQCD calculation, and for several event generators (dashed and dot-dashed lines). The gray band reflects uncertainties on the pQCD scale and parton distribution functions. Inner error bars represent the statistical uncertainty, whereas the outer one is statistical and systematic added in quadrature. b) shows the ratio of $(1/\sigma)d\sigma/dp_T$ relative to NLO pQCD for an approximate NNLO pQCD calculation and of predictions for several event generators.

$|\eta| < 1.1$. A cut on the missing transverse energy of 20 GeV is applied. Furthermore at least four jets are required with $p_T > 20$ GeV and $|\eta| < 2.0$. Finally at least one jet needs to be identified as a b -jet. The hadronic W decay is used to constrain the Jet Energy Scale (JES). $M_{t\bar{t}}$ is reconstructed by using the four-vectors of the b -tagged jet and the three remaining leading jets in the event, the lepton and the transverse components of the neutrino momentum, given by \cancel{E}_T .




Fig. 3. – (Color online) Panel a) shows the differential $t\bar{t}$ cross section (circles) as a function of $M_{t\bar{t}}$ [4] compared to the standard model expectation (line). The SM uncertainty (green band) reflects all systematic uncertainties, except for the luminosity uncertainty in each bin. Panel b) shows limits on the ratio κ/M_{Pl} for gravitons which decay to top quarks in the Randall-Sundrum model, where the mass of the first resonance is fixed at 600 GeV. Values of $\kappa/M_{Pl} > 0.16$ are excluded at the 95% confidence level.

Figure 3a) shows the differential $t\bar{t}$ cross section as a function of $M_{t\bar{t}}$ compared to the standard model expectation using the proton PDF CTEQ5L PDF with a top mass of 175 GeV. The SM uncertainty reflects all systematic uncertainties, except for the luminosity uncertainty in each bin. Especially the tail of $M_{t\bar{t}}$ is sensitive to broad enhancements as well as to narrow resonances, which is why the agreement between data and SM expectation has been evaluated. There is no indication of beyond standard model contributions to the differential cross section. The analysis also measured the inclusive total cross section for $t\bar{t}$ production to: $\sigma = 6.9 \pm 1.0(\text{stat.} + \text{JES})$ pb, which is in good agreement with latest theoretical predictions [5,6] as well as with the DØ result. Furthermore the distribution has been used to derive a limit on gravitons which decay to top quarks in the Randall-Sundrum model. The mass of the first resonance is fixed to 600 GeV and gravitons are modeled using MadEvent plus Pythia. Figure 3b) shows the derived limits, values of $\kappa/M_{Pl} > 0.16$ are excluded at the 95% confidence level.

4. – Conclusion

Two differential cross section measurements have been presented. The cross section as a function of the transverse momentum of the top quark by DØ [3] and as a function of the invariant mass of the $t\bar{t}$ system by CDF [4]. Both presented results are consistent with the standard model cross section predictions. The final Tevatron data sample has 5–10 times the presented statistics allowing for more precise measurements in the near future being one of the legacy measurements of the Tevatron.

REFERENCES

- [1] ABE F. *et al.* (CDF COLLABORATION), *Phys. Rev. Lett.*, **74** (1995) 2626 [[arXiv:hep-ex/9503002](#)].
- [2] ABACHI S. *et al.* (DØ COLLABORATION), *Phys. Rev. Lett.*, **74** (1995) 2632 [[arXiv:hep-ex/9503003](#)].
- [3] ABAZOV V. *et al.* (DØ COLLABORATION), *Phys. Lett. B*, **693** (2010) 515 [[arXiv.org:1001.1900](#)].
- [4] AALTONEN T. *et al.* (CDF COLLABORATION), *Phys. Rev. Lett.*, **102** (2003) 22 [[arXiv.org:0903.2850](#)].
- [5] AHRENS V., FERROGLIA A., NEUBERT M., PECJAK B. D. and YANG L. L., *JHEP*, **09** (2010) 097; AHRENS V., FERROGLIA A., NEUBERT M., PECJAK B. D. and YANG L. L., *Nucl. Phys. Proc. Suppl.*, **205-206** (2010) 48.
- [6] MOCH S. and UWER P., *Phys. Rev. D*, **78** (2008) 034003; LANGENFELD U., MOCH S. and UWER P., *Phys. Rev. D*, **80** (2009) 054009; ALIEV M. *et al.*, *Comput. Phys. Commun.*, **182** (2011) 1034.