

Parity violating neutron spin rotation in ^4He and H

W. M. SNOW⁽¹⁾, E. ANDERSON⁽¹⁾, L. BARRÓN-PALOS⁽²⁾, C. D. BASS⁽³⁾,
T. D. BASS⁽³⁾, B. E. CRAWFORD⁽⁴⁾, J. FRY⁽¹⁾, K. GAN⁽⁵⁾, B. R. HECKEL⁽⁶⁾,
D. LUO⁽¹⁾, R. C. MALONE⁽⁴⁾, D. M. MARKOFF⁽⁷⁾, A. M. MICHERDZINSKA⁽⁸⁾,
H. P. MUMM⁽⁹⁾, J. S. NICO⁽⁹⁾, A. K. OPPER⁽⁸⁾, S. PENN⁽¹⁰⁾, S. SANTRA⁽¹¹⁾,
M. SARSOOR⁽¹²⁾, E. I. SHARAPOV⁽¹³⁾, H. E. SWANSON⁽⁶⁾, S. VAN SCIVER⁽¹⁴⁾,
S. B. WALBRIDGE⁽¹⁾ and V. ZHUMABEKOVA⁽¹⁵⁾

⁽¹⁾ *Indiana University/CEEM - 2401 Milo B. Sampson Lane, Bloomington, IN 47408, USA*

⁽²⁾ *Universidad Nacional Autónoma de México - C.P. 04510, Mexico, D.F*

⁽³⁾ *Thomas Jefferson National Accelerator Facility - 12000 Jefferson Avenue
Newport News, VA 23606, USA*

⁽⁴⁾ *Gettysburg College - 300 North Washington Street, Gettysburg, PA 17325, USA*

⁽⁵⁾ *University of Manitoba - Winnipeg, MB, Canada R3T 2N2*

⁽⁶⁾ *University of Washington/CENPA - Box 354290, Seattle, WA 98195, USA*

⁽⁷⁾ *North Carolina Central University/TUNL - 1801 Fayetteville Street
Durham, NC 27707, USA*

⁽⁸⁾ *The George Washington University - 725 21st Street NW, Washington, D.C. 20052, USA*

⁽⁹⁾ *National Institute of Standards and Technology - 100 Bureau Drive
Gaithersburg, MD 20899, USA*

⁽¹⁰⁾ *Hobart William Smith College - Geneva, NY 14456, USA*

⁽¹¹⁾ *Bhabha Atomic Research Center - Trombay, Mumbai 400085, India*

⁽¹²⁾ *Georgia State University - 29 Peachtree Center Avenue, Atlanta, GA 30303-4106, USA*

⁽¹³⁾ *Joint Institute for Nuclear Research - Joliot-Curie 6, 141980 Dubna, Russia*

⁽¹⁴⁾ *National High Magnetic Field Laboratory and Florida State University
1800 E. Paul Dirac, Dr. Tallahassee, FL 32310-3706, USA*

⁽¹⁵⁾ *Al-Farabi Kazakh National University - Al-Farabi Ave. 71, 050038 Almaty, Kazakhstan*

ricevuto il 14 Ottobre 2011; approvato il 5 Maggio 2012
pubblicato online il 4 Luglio 2012

Summary. — The weak interaction between nucleons leads to parity violation in various reaction observables. Neutron spin rotation, the rotation of the plane of polarization of a transversely polarized neutron beam passing through unpolarized matter, is an especially clear example of a breakdown in mirror symmetry. The Neutron Spin Rotation (NSR) Collaboration is engaged in an experimental program to observe parity-odd neutron spin rotation. We recently completed the first phase of an experiment to measure parity violating neutron spin rotation in ^4He . Our result for the neutron spin rotation angle per unit length in ^4He , $d\phi/dz = (+1.7 \pm 9.1(\text{stat.}) \pm 1.4(\text{sys.})) \times 10^{-7} \text{ rad/m}$, is the most sensitive search for neutron weak optical activity yet performed and represents a significant advance in precision in comparison to past measurements in heavy nuclei. This experiment was performed at the NG-6 slow neutron beamline at the National Institute of Standards and Technology (NIST) Center for Neutron Research. The systematic uncertainty is small enough to proceed to the second phase of the ^4He measurement at the new NG-C slow neutron beamline under construction at NIST. The projected intensity of this beam is high enough to see parity odd neutron spin rotation in ^4He and to seriously consider a future experiment to measure neutron spin rotation in hydrogen.

PACS 11.30.Er – Charge conjugation, parity, time reversal, and other discrete symmetries.

PACS 24.70.+s – Polarization phenomena in reactions.

PACS 13.75.Cs – Nucleon-nucleon interactions (including antinucleons, deuterons, etc.).

1. – Introduction and theoretical overview

Neutron spin rotation is expected from quark-quark weak interactions in the Standard Model, which induce weak interactions between nucleons that violate parity. Because the range for W and Z exchange between quarks is small compared to the nucleon size, NN weak interaction amplitudes are one of the few observables which are first-order sensitive to quark-quark correlations in the nucleon. At energies well below the electroweak scale, the quark-quark weak interaction can be written in a current-current form with pieces that transform under isospin as $\Delta I = 0, 1, 2$. Lattice gauge theory [1] can be used to calculate the long-range component of the $\Delta I = 1$ NN weak interaction, and the first such calculation has recently been completed [2]. NN weak interactions also induce parity-odd effects in atomic structure, where they are the microscopic source for nuclear anapole moments [3-6].

Parity-odd neutron spin rotation has been measured in heavy nuclei [7-9], but the dynamics are too complicated to use this information to learn about the NN weak interaction amplitudes. To do this one must measure parity-odd neutron spin rotation and other parity-odd observables in light nuclei such as H, D, ^3He , and ^4He . Because strong interaction effects are now calculable [10] in few-body nuclei and weak amplitudes can be added as a perturbation, several new calculations of parity-odd effects in these systems have appeared recently [11-17] to complement earlier works [18-20]. The expected size of the parity-odd rotation angle in such few-body systems is about 10^{-6} rad/m to 10^{-7} rad/m [21].




Fig. 1. – Overview of an apparatus to measure parity-odd neutron spin rotation in liquid helium.

Of the five independent weak transition amplitudes present in NN elastic scattering at low energy [22], only the 1S_0 - 3P_0 proton-proton amplitude is fixed from experiment [23-26]; the rest are unknown. The existing calculation of $d\phi/dz$ in n - ${}^4\text{He}$ ($d\phi/dz = -0.97f_\pi - 0.22h_\omega^0 + 0.22h_\omega^1 - 0.32h_\rho^0 + 0.11h_\rho^1$) [27] was conducted within the meson exchange picture developed by Desplanques, Donoghue, and Holstein (DDH) [18] which uses π , ρ , and ω exchange parametrized by weak couplings at the NN vertex labeled by superscripts which indicate the isospin change. Within the DDH approach $d\phi/dz$ in n - ${}^4\text{He}$ spans a range of $\pm 1.5 \times 10^{-6}$ rad/m: this broad range of possibilities is dominated by the uncertainties in the weak couplings and reflects in part our poor understanding of quark-quark correlation physics in QCD. $d\phi/dz$ in n - ${}^4\text{He}$ has been related to existing measurements of nuclear parity violation in a model [20] which subsumes many poorly-understood short-range NN effects by expressing parity-odd amplitudes in terms of isoscalar ($X_n + X_p$) and isovector ($X_n - X_p$) one-body potentials. n - ${}^4\text{He}$ spin rotation is interesting within the context of this model since it determines X_n . Within this model measurements in ${}^{18}\text{F}$ [28] constrain $X_n - X_p$ and measurements in odd-proton systems such as p - ${}^4\text{He}$ [29,30] and ${}^{19}\text{F}$ [31] constrain X_p . The prediction in this model for n - ${}^4\text{He}$ spin rotation is $d\phi/dz = (-6.5 + / - 2.2) \times 10^{-7}$ rad/m.

2. – Neutron spin rotation and experimental technique

The phenomenon of neutron spin rotation can be understood in terms of neutron optics. The parity-violating (PV) weak interaction between the neutrons and the medium causes the amplitudes of the positive and negative neutron helicity states of polarized neutrons to accumulate different phases. The difference ϕ_{PV} between the phase shifts of the helicity states leads to a rotation of the neutron polarization vector about its momentum, which manifestly violates parity [32]. The rotation angle per unit length of a neutron of wave vector k in a medium of density ρ is $d\phi/dz = 4\pi\rho f_{PV}/k$, where f_{PV} is the forward limit of the parity-odd p -wave scattering amplitude. Because f_{PV} is proportional to the parity-odd correlation $\vec{\sigma}_n \cdot \vec{k}_n$ with $\vec{\sigma}_n$ the neutron spin vector and \vec{k}_n the neutron momentum, $d\phi/dz$ is constant as $k \rightarrow 0$ in the absence of resonances [33].

The experimental technique has been presented in detail [34,35]. The apparatus shown in fig. 1 must distinguish small PV rotations from rotations that arise from magnetic fields. ϕ_{PV} is isolated by alternately moving the medium in front of and behind a neutron spin precession coil and measuring the change in the spin rotation angle using the neutron equivalent of a crossed polarizer/analyzer pair familiar from light optics.

Neutrons polarized along \hat{y} enter a central precession coil with an internal magnetic field along \hat{y} (π -coil) which precesses a spin component along $+\hat{x}$ to $-\hat{x}$. The contribution to the total rotation angle coming from parity violation in the liquid changes sign as the liquid is moved. To further suppress systematic uncertainties and noise, the beam and apparatus are split into right and left halves, and the targets are filled so that the liquid occupies the chamber downstream of the π -coil on one side and the chamber upstream of the π -coil on the other side. The PV components of the neutron spin rotation angle have opposite signs on each side, and the difference of the two rotation angles is insensitive to both static residual magnetic fields and any common-mode time-dependent magnetic field integrals along the neutron trajectories.

The first phase of the measurement of neutron spin rotation in ^4He was performed at the NG-6 slow neutron beamline at the National Institute of Standards and Technology (NIST) Center for Neutron Research [36]. The neutrons were polarized vertically by a polarizing supermirror [37] and enter the magnetic shield/target region using a glass neutron guide and a magnetic field from an input coil to transport and preserve the neutron polarization [38]. The target vessel was mounted inside a magnetic shield centered in a nonmagnetic liquid helium cryostat and supported in turn inside two more layers of magnetic shielding. The liquid is moved between the four separate target chambers using a centrifugal pump immersed in a 4 K liquid helium bath outside the target with flexible tubes pulled by strings to determine which pair of target chambers fill or drain [34]. Internal fluxgate magnetometers indicate a typical internal axial magnetic field of 10 nT. After the target region an output coil and another float glass neutron guide conducts the transmitted neutrons to the polarization analyzer. The output coil adiabatically rotates the x -component of the neutron polarization by $\pm\pi/2$ in the x - y plane through modulation of the current direction in one of two orthogonal solenoids. This rotated spin component points along $\pm\hat{y}$ at the analyzer position to produce an asymmetry in the flux transmitted through the polarization analyzer given by $\frac{N_+ - N_-}{N_+ + N_-} = \langle PA \sin \phi \rangle$, where PA is the product of the neutron polarization from the polarizer and the analyzing power of the analyzer, ϕ is the neutron spin rotation angle, and N_+ and N_- are the count rates for $+$ and $-$ states of the output coil. The average is taken over the neutron velocity spectrum of the beam. The ion chamber operates in current mode using the $n + ^3\text{He} \rightarrow ^3\text{H} + p$ reaction and possesses four charge-sensitive collection plates along the beam direction with each plate subdivided into four quadrants [39]. The parity-odd spin rotation angle is constructed from the angles measured in the left and right target chambers and in the $+$ and $-$ π -coil states. A “ π -off” rotation angle is constructed from the angles measured with no current in the π -coil; this asymmetry must give zero in the absence of systematic errors. Possible false effects from slow drifts in the polarimetry were suppressed by analyzing the time sequence of asymmetries with an algorithm which cancels linear and quadratic time-dependent effects [40]. The left-right beam and target segmentation was essential to suppress common-mode nonstatistical noise from reactor power fluctuations in the 1 Hz frequency bandwidth of the rotation angle measurements: the noise was reduced by a factor of 8 [34]. The statistical uncertainty from the distribution of asymmetries is $\approx 15\%$ larger than would be expected from neutron counting statistics; about 8% of this extra noise comes from magnetic field fluctuations not removed by the filtering algorithm and a few percent comes from the current-mode operation of the ion chamber. The measured π -coil off angle of $(-1.2 \pm 10.0(\text{stat.})) \times 10^{-7}$ rad/m which is interleaved between the π -coil on measurements places an upper bound on the sum of all systematic effects. It is about a factor of 4 more sensitive than the π -on data to any systematic effects coupled to a constant longitudinal magnetic field but is not sufficiently precise to reduce the systematic

uncertainty to the required level. This was done in separate measurements and calculations [41]. Our result for the neutron spin rotation angle per unit length in ${}^4\text{He}$ in the first phase of the measurement, $d\phi/dz = (+1.7 \pm 9.1(\text{stat.}) \pm 1.4(\text{sys.})) \times 10^{-7}$ rad/m [42], is consistent with zero and with a previous unpublished result [43].

3. – Future measurements of neutron spin rotation in helium and hydrogen

The small size of the systematic uncertainty and the good prospects for stronger suppression of the internal magnetic fields encourage a next attempt in liquid ${}^4\text{He}$ on a more intense slow neutron beam. The new NG-C slow neutron beam under construction at NIST [44] coupled with a new supermirror polarizer is projected to provide a neutron fluence of 7×10^{10} n/s, a factor of 60 greater than that used in the previous experiment. We plan to improve the apparatus by using better-optimized magnetic shielding and control of external field fluctuations, an improved liquid helium pump and a helium liquifier to reduce deadtime from liquid motion and cryogen transfers, a neutron polarizer and analyzer of improved phase space uniformity, and supermirror input and output guides. We expect to reduce the statistical uncertainty on $d\phi/dz$ to 2×10^{-7} rad/m with smaller systematic uncertainties. This precision would place a tight constraint on a linear combination of NN weak amplitudes.

More theoretical work promises to impact the interpretation of this measurement. A calculation of $d\phi/dz$ in n- ${}^4\text{He}$ using Greens Function Monte Carlo techniques now in progress [45] should greatly improve the precision of the relative weighting with which the different weak amplitudes contribute. Theoretical approaches based on effective field theory [11, 13, 14] are under construction which incorporate the chiral symmetry of QCD. In particular, a new calculation of neutron spin rotation in hydrogen and deuterium in pionless effective field theory has recently appeared [46].

Liquid hydrogen in the para molecular state (which has $J = L = S = 0$) does not depolarize a slow neutron beam, whose kinetic energies are well below the 15 meV threshold for the excitation of the para to ortho molecular transition, and the slow neutron scattering cross section in parahydrogen is small enough that a target length of about 20 cm can be used. Both the successful operation of the liquid parahydrogen target for the NPDGamma experiment at LANSCE [47] and the successful use of slow neutron transmission to measure its (very small) orthohydrogen content [48] bode well for the development of a liquid parahydrogen target that could be used to measure parity odd neutron spin rotation in liquid parahydrogen.

* * *

This work is supported in part by the NSF PHY-0758018 and DOE DE-FG02-95ER40901 (USA), SEP-CONACYT (Mexico), and BARC (India). We acknowledge the support of the National Institute of Standards and Technology, US Department of Commerce, in providing the neutron facilities used in this work. WMS acknowledges support from the Indiana University Center for Spacetime Symmetries.

REFERENCES

- [1] BEANE S. R. and SAVAGE M. J., *Nucl. Phys. B*, **636** (2002) 291.
- [2] WASEM J., arXiv:1108.1151v1 [hep-lat] (2011).
- [3] ZELDOVICH Y. B., *Sov. Phys. JETP*, **6** (1957) 1184.
- [4] FLAMBAUM V. V. and KHRIPLOVICH I. B., *Sov. Phys. JETP*, **52** (1980) 835.

- [5] WOOD C. S. *et al.*, *Science*, **275** (1997) 1759.
- [6] TSIGUTKIN K. *et al.*, *Phys. Rev. Lett.*, **103** (2009) 071601.
- [7] FORTE M. *et al.*, *Phys. Rev. Lett.*, **45** (1980) 2088.
- [8] HECKEL B. R. *et al.*, *Phys. Lett. B*, **119** (1982) 298.
- [9] HECKEL B. R. *et al.*, *Phys. Rev. C*, **29** (1984) 2389.
- [10] PIEPER S. C. and WIRINGA R. B., *Annu. Rev. Nucl. Part. Sci.*, **51** (2001) 53.
- [11] ZHU S. L. *et al.*, *Nucl. Phys. A*, **748** (2005) 435.
- [12] RAMSEY-MUSOLF M. J. and PAGE S., *Annu. Rev. Nucl. Part. Sci.*, **56** (2006) 2.
- [13] LIU C. P., *Phys. Rev. C*, **75** (2007) 066501.
- [14] PHILLIPS D. R., SCHINDLER M. R. and SPRINGER R. P., *Nucl. Phys. A*, **822** (2009) 1.
- [15] SCHIAVILLA R., VIVANI M., GIRLANDA L., KIEVSKY A. and MARCUCCI L. E., *Phys. Rev. C*, **78** (2008) 014002.
- [16] HWANG W.-Y. P. and WEN C. Y., *Phys. Rev. C*, **78** (2008) 022501.
- [17] SHIN J. W., ANDO S. and HYUN C. H., *Phys. Rev. C*, **81** (2010) 055501.
- [18] DESPLANQUES B., DONOGHUE J. F. and HOLSTEIN B. R., *Ann. Phys. (N.Y.)*, **124** (1980) 449.
- [19] ADELBERGER E. G. and HAXTON W. C., *Annu. Rev. Nucl. Part. Sci.*, **35** (1985) 501.
- [20] DESPLANQUES B., *Phys. Rep.*, **297** (1998) 1.
- [21] STODOLSKY L., *Phys. Lett. B*, **50** (1974) 353.
- [22] DANILOV G. S., *Phys. Lett.*, **18** (1965) 40.
- [23] POTTER J. M. *et al.*, *Phys. Rev. Lett.*, **33** (1974) 1307.
- [24] BALZER R. *et al.*, *Phys. Rev. Lett.*, **44** (1980) 699.
- [25] KISTRYN S. *et al.*, *Phys. Rev. Lett.*, **58** (1987) 1616.
- [26] EVERSHEIM P. D. *et al.*, *Phys. Lett. B*, **256** (1991) 11.
- [27] DMITRIEV V. *et al.*, *Phys. Lett.*, **125** (1983) 1.
- [28] BARNES C. A. *et al.*, *Phys. Rev. Lett.*, **40** (1978) 840; PAGE S. A. *et al.*, *Phys. Rev. C*, **35** (1987) 1119; EVANS H. C. *et al.*, *Phys. Rev. Lett.*, **55** (1985) 791; BINI M. *et al.*, *Phys. Rev. Lett.*, **55** (1985) 795.
- [29] LANG J. *et al.*, *Phys. Rev. Lett.*, **54** (1985) 170.
- [30] LANG J. *et al.*, *Phys. Rev. C*, **34** (1986) 1545.
- [31] ELSENER K. *et al.*, *Phys. Rev. Lett.*, **52** (1984) 1476.
- [32] MICHEL F. C., *Phys. Rev. B*, **133** (1964) 329.
- [33] STODOLSKY L., *Nucl. Phys. B*, **197** (1982) 213.
- [34] BASS C. D. *et al.*, *Nucl. Instrum. Methods Phys. Res. A*, **612** (2009) 69.
- [35] SNOW W. M., *Nucl. Instrum. Methods Phys. Res. A*, **611** (2009) 248.
- [36] NICO J. S. *et al.*, *J. Res. Natl. Inst. Stand. Technol.*, **110** (2005) 137.
- [37] SCHAERPF O., *Physica B*, **156-157** (1989) 631; (1989) 639.
- [38] FORTE M., in *Physics with Reactor Neutrons and Neutrinos*, edited by VON EGIDY T., Vol. **42** (Institute of Physics Conference Series) 1978, p. 80.
- [39] PENN S. *et al.*, *Nucl. Instrum. Methods Phys. Res. A*, **457** (2001) 332.
- [40] SWANSON H. E. and SCHLAMMINGER S., *Meas. Sci. Technol.*, **21** (2010) 115104.
- [41] MICHARDZINSKA A. *et al.*, *Nucl. Instrum. Methods Phys. Res. A*, **631** (2011) 809.
- [42] SNOW W. M. *et al.*, *Phys. Rev. C*, **83** (2011) 022501(R).
- [43] MARKOFF D., PhD thesis, University of Washington (1997).
- [44] COOK J. C., *Rev. Sci. Instrum.*, **80** (2009) 023101.
- [45] SCHIAVILLA R., private communication (2010).
- [46] GRIESSHAMMER H. W., SCHINDLER M. and SPRINGER R. P., arXiv:1109.5667v1 [nucl-th] (2011).
- [47] SANTRA S. *et al.*, *Nucl. Instrum. Methods Phys. Res. A*, **620** (2010) 421.
- [48] BARRON-PALOS L. *et al.*, *Nucl. Instrum. Methods Phys. Res. A*, **659** (2011) 579.