

Kaon-nuclei interaction studies at low energies (the AMADEUS project)

K. PISCICCHIA^{(1)(2)(*)}, M. BAZZI⁽¹⁾, C. BERUCCI⁽¹⁾, D. BOSNAR⁽³⁾,
A. M. BRAGADIREANU⁽⁴⁾, M. CARGNELLI⁽⁵⁾, A. CLOZZA⁽¹⁾, C. CURCEANU⁽¹⁾,
A. D'UFFIZI⁽¹⁾, F. GHIO⁽⁶⁾, C. GUARALDO⁽¹⁾, P. KIENLE⁽⁵⁾⁽⁷⁾, M. ILIESCU⁽¹⁾,
T. ISHIWATARI⁽⁵⁾, P. LEVI SANDRI⁽¹⁾, J. MARTON⁽⁵⁾, D. PIETREANU⁽¹⁾⁽⁴⁾,
M. POLI LENER⁽¹⁾, A. RIZZO⁽¹⁾, A. ROMERO VIDAL⁽¹⁾, E. SBARDELLA⁽¹⁾,
A. SCORDO⁽¹⁾, D. L. SIRGHI⁽¹⁾⁽⁴⁾, F. SIRGHI⁽¹⁾⁽⁴⁾, H. TATSUNO⁽¹⁾, I. TUCAKOVIC⁽¹⁾,
O. VAZQUEZ DOCE⁽⁸⁾, E. WIDMANN⁽⁵⁾ and J. ZMESKAL⁽⁵⁾

⁽¹⁾ INFN, Laboratori Nazionali di Frascati - Frascati (RM), Italy

⁽²⁾ Dipartimento di Fisica, Università "Roma Tre" - Roma, Italy

⁽³⁾ Physics Department, University of Zagreb - Zagreb, Croatia

⁽⁴⁾ IFIN-HH, Mağurele - Bucharest, Romania

⁽⁵⁾ Stefan-Meyer-Institut für subatomare Physik - Vienna, Austria

⁽⁶⁾ INFN, Sezione di Roma I and Istituto Superiore di Sanità - Roma, Italy

⁽⁷⁾ Technische Universität München, Physik Department - Garching, Germany

⁽⁸⁾ Excellence Cluster Universe, Technische Universität München - Boltzmannstr. 2
D-85748, Garching, Germany

ricevuto il 31 Agosto 2012

Summary. — The AMADEUS experiment aims to perform dedicated precision studies in the sector of low-energy kaon-nuclei interaction at the DAΦNE collider at LNF-INFN. In particular the experiment plans to perform measurements of the debated deeply bound kaonic nuclear states (by stopping kaons in cryogenic gaseous targets ³He and ⁴He) to explore the nature of the $\Lambda(1405)$ in nuclear environment and to measure the cross section of K^- on light nuclei, for K^- momentum lower than 100 MeV/c. The AMADEUS dedicated setup will be installed in the central region of the KLOE detector

PACS 13.75.Jz – Kaon-baryon interactions.

PACS 25.80.Nv – Kaon-induced interactions.

PACS 21.65.Jk – Mesons in nuclear matter.

(*) E-mail: kristian.piscicchia@lnf.infn.it

1. – Introduction

The AMADEUS (Antikaon Matter At DAΦNE Experiments with Unraveling Spectroscopy) experiment [1, 2] will study the low-energy interactions of kaons with nucleons and nuclei. AMADEUS aims to search for the most fundamental Deeply Bound Kaonic Nuclear States (DBKNS), that are the kaonic dibaryon states (K^-pp , K^-pn), produced by stopping K^- in a ^3He target and, as a next step, the kaonic tribaryon states (K^-ppn , K^-pnn) using a ^4He target. A great experimental effort is ongoing in the search for such states, but the possible experimental indications of the formation of dibaryonic and tribaryonic states, have received alternative explanations in the framework of known processes [3]. Another important subject is the $\Lambda(1405)$ resonance [4], whose nature is still not completely understood. The structure of the $\Lambda(1405)$ has been found to be important in various aspects in the strangeness sector of nonperturbative QCD. AMADEUS will give the possibility to better understand such state and its behavior in nuclear environment, with high statistics. Moreover, AMADEUS plans to perform the measurement of low energy cross sections of charged kaons on H, d, and He (for kaons momentum lower than $100\text{ MeV}/c$) and the study of nuclear interactions of K^- in various targets.

2. – Setup performance requirements

Kaonic nuclear states will be studied both in the formation process, by means of missing mass spectroscopy, and in the decay process. As DBKNS are expected to decay into states containing Λ and Σ hyperons, neutral (neutrons and photons) as well as charged particles are to be detected, with momenta in a wide range. All these detection requirements are perfectly satisfied by the KLOE detector, which is made of a 4π cylindrical drift chamber (DC) and a calorimeter, both immersed in the 0.52 T field of a superconducting solenoid. KLOE has an acceptance of 96%, is optimized for detection of charged particles in the relevant energy range, and has good detection efficiency for neutrons, as was checked by the KloNe group [5]. On the other side, DAΦNE is a unique source of low energy kaons. DAΦNE is a double ring e^+e^- collider, designed to work in the center of mass energy of the ϕ meson. Charged kaons coming from ϕ decay are characterized by low momentum ($\sim 127\text{ MeV}/c$), which enables to stop them in gaseous targets, and a back to back topology which turns to be ideal for triggering purpose.

3. – The dedicated AMADEUS setup

The AMADEUS setup will be implemented inside the KLOE DC, between the beam pipe (6 cm diameter) and the DC entrance wall (50 cm diameter). Two main components of the experimental setup are presently under development: a high density cryogenic gaseous target and a trigger system.

An essential feature of the detector is the possibility to trigger on charged kaons coming from the interaction point. The main goal is a time resolution sufficient to clearly distinguish kaons from background. This will be achieved by making use of two layers of scintillating fibers read at both sides by silicon photomultipliers (SiPM) surrounding the beam pipe. Employing two high granularity layers will give the possibility to perform a preliminary tracking as well. SiPM turn to be optimal for our purposes as they are rather insensitive to magnetic field and are characterized by reduced dimensions. A prototype of the SiPM + SciFi system was already tested on DAΦNE (fibers were placed under the lower scintillator of the SIDDHARTA Kaon Monitor) [6, 7]. A second and more complex prototype, constituted of two layers of BCF-10 double cladded fibers, free

to rotate and read at both sides by Hamamatsu S10362-11-050-U SiPMs was recently tested. An excellent time resolution for kaons was achieved ($\sigma \sim 300$ ps) [8].

A half toroidal cryogenic target is under study, enclosed in a vacuum chamber, filled with ^3He as a first step, ^4He in a second phase. According to MC simulation about 20% of the negative kaons from DAΦNE should be stopped in the gas filling the target. More outer layers of scintillating fibers opposite to the target cell would enable to clearly identify the K^+ , and to perform a reconstruction of the inner trajectory of the kaons. A similar target was recently installed in DAΦNE, for the SIDDHARTA [9-11] experiment and our group will take advantage of the gained experience.

4. – Analysis of the KLOE data searching for K^- - ^4He interactions

Presently, we are performing dedicated Monte Carlo simulations, to study the performance of the AMADEUS setup. In parallel, we are analyzing the existing KLOE data (runs from 2002 to 2005). Indeed the KLOE drift chamber is mainly filled with ^4He (90% helium 10% isobutane) and the analysis of KLOE Monte Carlo showed, that about 0.1% of kaons from DAΦNE should stop in the inner volume of the drift chamber. This represents a unique opportunity to check the reconstruction capability for Σ and Λ particles and for studying the hadronic interactions of K^- in such an active target. Up to now data for a total luminosity of 1.8 fb^{-1} were analyzed [12]. An excellent result was already achieved in reconstructing the $\Lambda(1116)$ invariant mass, with a statistical error $\sigma \sim 3\text{ keV}$; the systematics being presently under evaluation. The investigation of the Λp and Λd events, together with the study of the $\Lambda(1405)$ through the neutral decay channel $\Sigma^0\pi^0$ already gave excellent results, which are presently being finalized.

* * *

Part of this work was supported by HadronPhysics I3 FP6 European Community program, Contract No. RII3-CT-2004-506078; the European Community-Research Infrastructure Integrating Activity “Study of Strongly Interacting Matter” (HadronPhysics 2, Grant Agreement No. 227431), and HadronPhysics 3 (HP3), Contract No. 283286 under the Seventh Framework Programme of EU.

REFERENCES

- [1] AMADEUS Letter of Intent, http://www.lnf.infn.it/esperimenti/siddharta/LOI-AMDEUS_March2006.pdf.
- [2] THE AMADEUS COLLABORATION, LNF preprint, LNF07/24(IR) (2007).
- [3] OSET E. and TOKI H., *Phys. Rev. C*, **74** (2006) 015207 [nucl-th/0509048].
- [4] HYODO TETSUO and JIDO DAISUKE, *Prog. Particle Nucl. Phys.*, **67** (2012) 55.
- [5] ANELLI M. *et al.*, *Nucl. Instrum. Methods A*, **581** (2007) 368.
- [6] BAZZI M., CORRADI G., PISCICCHIA K., ROMERO VIDAL A., SCORDO A., TAGNANI D. and VASQUEZ DOCE O., AMADEUS Technical Note IR-2 04/10/2009.
- [7] BAZZI M., CORRADI G., PISCICCHIA K., ROMERO VIDAL A., SCORDO A., TAGNANI D. and VASQUEZ DOCE O., AMADEUS Technical Note IR-1 29/09/2009.
- [8] BAZZI M., BERUCCI C., CORRADI G., CURCEANU C., D’UFFIZI A., PISCICCHIA K., POLI LENER M., RIZZO A., ROMERO VIDAL A., SBARDELLA E., SCORDO A., TAGNANI D. and VASQUEZ DOCE O., *Nucl. Instrum. Methods Phys. Res. A*, **671** (2012) 1251.
- [9] BAZZI M. *et al.*, *Phys. Lett. B*, **697** (2011) 199.
- [10] BAZZI M. *et al.*, *Phys. Lett. B*, **681** (2009) 310.
- [11] BAZZI M. *et al.*, *Phys. Lett. B*, **704** (2011) 113.
- [12] CARGNELLI M., PETRASCU C., VASQUEZ DOCE O. and KLOE K-CHARGED GROUP, KLOE memo 337 (2007).