

Search for dark matter in the sky with the Fermi Large Area Telescope

A. MORSELLI

INFN Roma Tor Vergata - Roma, Italy

ricevuto il 20 Giugno 2013; approvato l'1 Luglio 2013

Summary. — Can we learn about New Physics with astronomical and astroparticle data? Since its launch in 2008, the Large Area Telescope, onboard of the Fermi Gamma-ray Space Telescope, has detected the largest amount of gamma rays in the 20 MeV–300 GeV energy range and electrons + positrons in the 7 GeV–1 TeV range. These impressive statistics allow one to perform a very sensitive indirect experimental search for dark matter. We will present the latest results on these searches and the comparison with LHC searches.

PACS 96.50.sb – Composition, energy spectra and interactions.

PACS 95.35.+d – Dark matter (stellar, interstellar, galactic, and cosmological).

PACS 95.85.Ry – Neutrino, muon, pion, and other elementary particles; cosmic rays.

PACS 98.70.Sa – Cosmic rays (including sources, origin, acceleration, and interactions).

1. – Introduction

The Fermi Observatory carries two instruments on-board: the Gamma-ray Burst Monitor (GBM) [1] and the Large Area Telescope (LAT) [2]. The LAT is a pair conversion telescope for photons above 20 MeV up to a few hundreds of GeV. The field of view is ~ 2.4 sr and LAT observes the entire sky every ~ 3 hours (2 orbits). These features make the LAT a great instrument for dark matter (DM) searches. The operation of the instrument through the first three years of the mission was smooth at a level which is probably beyond the more optimistic pre-launch expectations. The LAT has been collecting science data for more than 99% of the time spent outside the South Atlantic Anomaly (SAA). The remaining tiny fractional down-time accounts for both hardware issues and detector calibrations [3, 4].

More than 650 million gamma-ray candidates (*i.e.* events passing the background rejection selection) were made public and distributed to the Community through the Fermi Science Support Center (FSSC)⁽¹⁾.

⁽¹⁾ The FSSC is available at <http://fermi.gsfc.nasa.gov/ssc>




Fig. 1. – Sky map of the energy flux derived from 24 months of observation. The image shows γ -ray energy flux for energies between 100 MeV and 10 GeV, in units of 10^{-7} erg cm $^{-2}$ s $^{-1}$ sr $^{-1}$.

Over the first three years of mission the LAT Collaboration has put a considerable effort toward a better understanding of the instrument and of the environment in which it operates. In addition to that a continuous effort was made in order to make the advances public as soon as possible. In August 2011 the first new event classification (Pass 7) since launch was released, along with the corresponding Instrument Response Functions (and a release of a new event class “Pass 7 reprocessed” is planned for the near future). Compared with the pre-launch (Pass 6) classification, it features a greater and more uniform exposure, with a significance enhancement in acceptance below 100 MeV.

2. – The second Fermi-LAT catalog

The high-energy gamma-ray sky is dominated by diffuse emission: more than 70% of the photons detected by the LAT are produced in the interstellar space of our Galaxy by interactions of high-energy cosmic rays with matter and low-energy radiation fields. An additional diffuse component with an almost-isotropic distribution (and therefore thought to be extragalactic in origin) accounts for another significant fraction of the LAT photon sample. The rest consists of various different types of point-like or extended sources: Active Galactic Nuclei (AGN) and normal galaxies, pulsars and their relativistic wind nebulae, globular clusters, binary systems, shock-waves remaining from supernova explosions and nearby solar-system bodies like the Sun and the Moon.

The Second Fermi-LAT catalog (2FGL) [5] is the deepest catalog ever produced in the energy band between 100 MeV and 100 GeV. Compared to the First Fermi-LAT (1FGL) [6], it features several significant improvements: it is based on data from 24 (*vs.* 11) months of observation and makes use of the new Pass 7 event selection. The energy flux map is shown in fig. 1. It is interesting to note that 127 sources are firmly identified, based either on periodic variability (*e.g.* pulsars) or on spatial morphology or on correlated variability. In addition to that 1170 are reliably associated with sources known at other wavelengths, while 576 (*i.e.* 31% of the total number of entries in the catalog) are still unassociated. In addition, the first catalog of high energy sources [7] as well as the first SNR catalog are in preparation [8].

3. – Indirect dark matter searches

One of the major open issues in our understanding of the Universe is the existence of an extremely-weakly interacting form of matter, the Dark Matter (DM), supported by a wide range of observations including large scale structures, the cosmic microwave background and the isotopic abundances resulting from the primordial nucleosynthesis. Complementary to direct searches being carried out in underground facilities and at accelerators, the indirect search for DM is one of the main items in the broad Fermi Science menu. The word indirect denotes here the search for signatures of Weakly Interactive Massive Particle (WIMP) annihilation or decay processes through the final products (gamma-rays, electrons and positrons, antiprotons) of such processes. Among many other ground-based and space-borne instruments, the LAT plays a prominent role in this search through a variety of distinct search targets: gamma-ray lines, Galactic and isotropic diffuse gamma-ray emission, dwarf satellites, CR electrons and positrons.

3.1. Galactic center. – The Galactic center (GC) is expected to be the strongest source of γ -rays from DM annihilation, due to its coincidence with the cusped part of the DM halo density profile [9-11]. A preliminary analysis of the data, taken during the first 11 months of the Fermi satellite operations is presented in [12, 13].

The diffuse gamma-ray backgrounds and discrete sources, as we know them today, can account for the large majority of the detected gamma-ray emission from the Galactic Center. Nevertheless a residual emission is left, not accounted for by the above models [12, 13]. Improved modeling of the Galactic diffuse model as well as the potential contribution from other astrophysical sources (for instance unresolved point sources) could provide a better description of the data. Analyses are underway to investigate these possibilities.

3.2. Galactic halo. – In order to minimize uncertainties connected with the region of the Galactic Center, analysis [14] considered a region of interest consisting of two off-plane rectangles ($5^\circ \leq |b| \leq 15^\circ$ and $|l| \leq 80^\circ$) and searched for continuum emission from dark matter annihilation or decay in the smooth Galactic dark matter halo. They considered two approaches: a more conservative one in which limits were set on DM models assuming that all gamma ray emission in that region might come from dark matter (*i.e.* no astrophysical signal is modeled and subtracted). In a second approach, dark matter source and astrophysical emission was fit simultaneously to the data, marginalizing over several relevant parameters of the astrophysical emission. As no robust signal of DM emission is found, DM limits are set.

These limits are particularly strong on leptonic DM channels, which are hard to constrain in most other probes (notably in the analysis of the dwarf Galaxies, described below). This analysis strongly challenges DM interpretation of the positron rise, observed by PAMELA [15] and Fermi LAT [16, 17] (see fig. 2).

3.3. Dwarf galaxies. – Dwarf satellites of the Milky Way are among the cleanest targets for indirect dark matter searches in gamma-rays. They are systems with a very large mass/luminosity ratio (*i.e.* systems which are largely DM dominated). The LAT detected no significant emission from any of such systems and the upper limits on the γ -ray flux allowed us to put very stringent constraints on the parameter space of well motivated WIMP models [18].

A combined likelihood analysis of the 10 most promising dwarf galaxies, based on 24 months of data and pushing the limits below the thermal WIMP cross section for




Fig. 2. – Derived 95% CL upper limits on WIMP annihilation cross sections in the Milky Way halo, for the muon (left) and tau (right) annihilation channels.

low DM masses (below a few tens of GeV), has been recently performed [19]. The main advantages of the combined likelihood are that the analysis can be individually optimized and that combined limits are more robust under individual background fluctuations and under individual astrophysical modelling uncertainties than individual limits. The derived 95% CL upper limits on WIMP annihilation cross sections for different channels are shown in fig. 3 (left). The most generic cross section ($\sim 3 \cdot 10^{-26} \text{ cm}^3 \text{ s}^{-1}$ for a purely s-wave cross section) is plotted as a reference. These results are obtained for NFW profiles [20] but for cored dark matter profile the J-factors for most of the dSphs would either increase or not change much so these results includes J-factor uncertainties [19].

With the present data we are able to rule out large parts of the parameter space where the thermal relic density is below the observed cosmological dark matter density and WIMPs are dominantly produced non-thermally, *e.g.* in models where supersymmetry breaking occurs via anomaly mediation for the MSSM model, updated from [18]).

Future improvements (apart from increased amount of data) will include an improved event selection with a larger effective area and photon energy range, and the inclusion of more satellite galaxies. In fig. 3 (right) are shown the predicted upper limits in the hypothesis of 10 years of data instead of 2; 30 dSphs instead of ten (supposing that the new optical surveys will find new dSph); spatial extension analysis (source extension increases the signal region at high energy $E \geq 10 \text{ GeV}$, $M \geq 200 \text{ GeV}$).




Fig. 3. – Left: Derived 95% CL upper limits on WIMP annihilation cross sections for different channels. Right: Predicted 95% CL upper limits on WIMP annihilation cross sections in 10 years for $b\bar{b}$ channel.




Fig. 4. – Dark-matter annihilation 95% CL cross section upper limits into $\gamma\gamma$ for the Einasto profile for a circular region of interest (ROI) with a radius $R_{GC} = 16^\circ$ centered on the GC with $|b| < 5^\circ$ and $|l| > 6^\circ$ masked.

Other complementary limits were obtained with the search of possible anisotropies generated by the DM halo substructures [21], the search for dark matter satellites [22] and a search for high-energy cosmic-ray electrons from the Sun [23].

3.4. Gamma-ray lines. – A line at the WIMP mass, due to the 2γ production channel, could be observed as a feature in the astrophysical source spectrum [11]. Such an observation would be a “smoking gun” for WIMP DM as it is difficult to explain by a process other than WIMP annihilation or decay and the presence of a feature due to annihilation into γZ in addition would be even more convincing. No significant evidence of gamma-ray line(s) has been found in the first two years of data from 7 to 200 GeV [24] (see also [25]).

Recently, the claim of an indication of line emission in Fermi-LAT data [26, 27] has drawn considerable attention. Using an analysis technique similar to [25], but doubling the amount of data as well as optimizing the region of interest for signal over square-root of background, [26] found a (trial corrected) 3.2σ significant excess at a mass of ~ 130 GeV that, if interpreted as a signal would amount to a cross-section of about $\langle\sigma v\rangle \sim 10^{-27} \text{ cm}^3 \text{ s}^{-1}$.

The signal is found to be concentrated on the Galactic Centre with a spatial distribution consistent with an Einasto profile [28]. This is marginally compatible with the upper limit presented in [24].

In the analysis of the 4 year data the Fermi LAT team has improved over the two year paper in three important aspects: i) the search was performed in five regions of interest optimized for DM search under five different assumptions on the morphology of the DM signal, ii) new improved data set (pass 7 reprocessed) was used, as it corrects for loss in calorimeter light yield due to radiation damage during the four years of the Fermi mission and iii) point spread function (PDF) was improved by adding a 2nd dimension to the previously used triple Gaussian PDF model, leading to a so called “2D” PDF (such procedure is shown to increase the sensitivity to a line detection by 15%). In that analysis [29] no globally significant lines have been found and new limits to this DM annihilation channel were set (see fig. 4). In a close inspection of the 130 GeV feature




Fig. 5. – Positron fraction measured by the Fermi LAT and by other experiments [15,30,31]. The Fermi statistical uncertainty is shown with error bars and the total (statistical plus systematic uncertainty) is shown as a shaded band.

it was found that indeed there exist a 135 GeV signal at 4.01σ local significance, when a “1D” PSF and old data sets were used (consistently with what [26, 27] have found). However, the significance drops to 3.35σ (local, or $\leq 2\sigma$ global significance once trials factors are taken into account). In addition, a weaker signal is found at the same energy in the control sample (in the Earth limb), which might point to a systematics effect present in this data set. In order to examine this possibility weekly observations of the Limb are scheduled, and a better understanding of a nature of the excess in the control sample should be available soon.

A new version of the event-level reconstruction and analysis framework (called Pass 8) is foreseen soon from the Fermi-LAT Collaboration. With this new analysis software we should increase the efficiency of the instrument at high energy and have a data set based on independent event analysis thus gaining a better control of the systematic effects.

3.5. The Cosmic Ray Electron spectrum. – The experimental information available on the Cosmic Ray Electron (CRE) spectrum has been dramatically expanded with a high precision measurement of the electron spectrum from 7 GeV to 1 TeV by the Fermi LAT [16, 17]. The spectrum shows no prominent spectral features and it is significantly harder than that inferred from several previous experiments.

More recently the Fermi-LAT Collaboration provided a further, and stronger, evidence of the positron anomaly by providing direct measurement of the absolute e^+ and e^- spectra, and of their fraction, between 20 and 200 GeV using the Earth magnetic field (see fig. 5). A steady rising of the positron fraction was observed up to that energy in agreement with that found by PAMELA. In the same energy range, the e^- spectrum was fitted with a power-law with index $\gamma(e^-) = -3.19 \pm 0.07$ which is in agreement with what recently measured by PAMELA between 1 and 625 GeV [15]. Most importantly, Fermi-LAT measured, for the first time, the e^+ spectrum in the 20–200 GeV energy interval (see fig. 5). The e^+ spectrum is fitted by a power-law with index $\gamma(e^+) = -2.77 \pm 0.14$.

These measurements seems to rule out the standard scenario in which the bulk of electrons reaching the Earth in the GeV - TeV energy range are originated by Supernova Remnants (SNRs) and only a small fraction of secondary positrons and electrons comes

from the interaction of CR nuclei with the interstellar medium (ISM). An additional electron + positron component peaked at ~ 1 TeV seems necessary for a consistent description of all the available data sets. The temptation to claim the discovery of dark matter from detection of electrons from annihilation of dark matter particles is strong but there are competing astrophysical sources, such as pulsars, that can give a strong flux of primary positrons and electrons (see [32] and references therein). At energies between 100 GeV and 1 TeV the electron flux reaching the Earth may be the sum of an almost homogeneous and isotropic component produced by Galactic supernova remnants and the local contribution of a few pulsars with the latter expected to contribute more and more significantly as the energy increases. If a single nearby pulsar give the dominant contribution to the extra component a large anisotropy and a small bumpiness should be expected; if several pulsars contribute the opposite scenario is expected.

So far no positive detection of CRE anisotropy was reported by the Fermi-LAT Collaboration, but some stringent upper limits were published [33] the pulsar scenario is still compatible with these upper limits.

After the conference the AMS-02 Collaboration presented the result on the positron fraction [34] that confirm the positron ratio rise observed by PAMELA and Fermi and extend it up to 350 GeV.

Forthcoming experiments like AMS-02 and CALET are expected to reduce drastically the uncertainties on the propagation parameters by providing more accurate measurements of the spectra of the nuclear components of CR. Fermi-LAT and those experiments are also expected to provide more accurate measurements of the CRE spectrum and anisotropy looking for features which may give a clue of the nature of the extra component.

4. – Conclusions

Fermi turned four years in orbit on June, 2012, and it is definitely living up to its expectations in terms of scientific results delivered to the community. The mission is planned to continue at least four more years (likely more) with many remaining opportunities for discoveries.

* * *

The Fermi-LAT Collaboration acknowledges support from a number of agencies and institutes for both development and the operation of the LAT as well as scientific data analysis. These include NASA and DOE in the United States, CEA/Irfu and IN2P3/CNRS in France, ASI and INFN in Italy, MEXT, KEK, and JAXA in Japan, and the K. A. Wallenberg Foundation, the Swedish Research Council and the National Space Board in Sweden. Additional support from INAF in Italy and CNES in France for science analysis during the operations phase is also gratefully acknowledged.

REFERENCES

- [1] MEEGAN C. *et al.*, *Astrophys. J.*, **702** (2009) 791.
- [2] ATWOOD W. B. *et al.* (FERMI COLLABORATION), *Astrophys. J.*, **697** (2009) 1071 arXiv:0902.1089.
- [3] ACKERMANN M. *et al.* (FERMI COLLABORATION), *Astropart. Phys.*, **35** (2012) 346353 arXiv:1108.0201.

- [4] ACKERMANN M. *et al.* (FERMI COLLABORATION), *Astrophys. J. Suppl. Ser.*, **203** (2012) 4 arXiv:1206.1896.
- [5] ABDO A. *et al.* (FERMI COLLABORATION), *Astrophys. J. Suppl. Ser.*, **199** (2012) 31 arXiv:1108.1435.
- [6] ABDO A. *et al.* (FERMI COLLABORATION), *Astrophys. J. Suppl. Ser.*, **188** (2010) 405 arXiv:1002.2280.
- [7] PANEQUE D., BALLEST J., BURNETT T., DIGEL S., FORTIN P. and KNOEDLSEDER J., arXiv:1304.4153 [astro-ph.HE].
- [8] DE PALMA F. *et al.* (FOR THE FERMI LAT COLLABORATION), arXiv:1304.1395 [astro-ph.HE].
- [9] MORSELLI A. *et al.*, *Nucl. Phys.*, **113B** (2002) 213.
- [10] CESARINI A., FUCITO F., LIONETTO A., MORSELLI A. and ULLIO P., *Astropart. Phys.*, **21** (2004) 267 astro-ph/0305075.
- [11] BALTZ E. *et al.*, *JCAP*, **07** (2008) 013 arXiv:0806.2911.
- [12] VITALE V. and MORSELLI A., for the Fermi/LAT Collaboration, 2009 Fermi Symposium arXiv:0912.3828.
- [13] MORSELLI A., CAÑADAS B. and VITALE V., *Nuovo Cimento C*, **34** (2011) 311 arXiv:1012.2292.
- [14] ACKERMANN M. *et al.* (FERMI COLLABORATION), *Astrophys. J.*, **761** (2012) 91 arXiv:1205.6474.
- [15] ADRIANI O. *et al.* (PAMELA COLLABORATION), *Phys. Rev. Lett.*, **106** (2011) 201101.
- [16] ABDO A. A. *et al.* (FERMI COLLABORATION), *Phys. Rev. Lett.*, **102** (2009) 181101 arXiv:0905.0025.
- [17] ACKERMANN M. *et al.* (FERMI COLLABORATION), *Phys. Rev. D*, **82** (2010) 092004 arXiv:1008.3999.
- [18] ABDO A. *et al.* (FERMI COLLABORATION), *Astrophys. J.*, **712** (2010) 147 arXiv:1001.4531.
- [19] ACKERMANN M. *et al.* (FERMI COLLABORATION), *Phys. Rev. Lett.*, **107** (2011) 241302 arXiv:1108.3546.
- [20] NAVARRO J., FRENK J. and WHITE S., *Astrophys. J.*, **462** (1996) 563 arXiv:astro-ph/9508025.
- [21] ACKERMANN M. *et al.* (FERMI COLLABORATION), *Phys. Rev. D*, **85** (2012) 083007 arXiv:1202.2856.
- [22] ACKERMANN M. *et al.* (FERMI COLLABORATION), *Astrophys. J.*, **747** (2012) 121 arXiv:1201.2691.
- [23] AJELLO M. *et al.* (FERMI COLLABORATION), *Phys. Rev. D*, **84** (2011) 032007 arXiv:1107.4272.
- [24] ACKERMANN M. *et al.* (FERMI COLLABORATION), *Phys. Rev. D*, **86** (2012) 022002 arXiv:1205.2739.
- [25] ABDO A. *et al.* (FERMI COLLABORATION), *Phys. Rev. Lett.*, **104** (2010) 091302 arXiv:1001.4836.
- [26] WENIGER C., *JCAP*, **08** (2012) 007 arXiv:1204.2797 [hep-ph].
- [27] SU M. and FINKBEINER D. P., arXiv:1206.1616 [astro-ph.HE].
- [28] BRINGMANN T. and WENIGER C., *Dark Universe*, **1** (2012) 194 arXiv:1208.5481.
- [29] BLOOM E. *et al.* (ON BEHALF OF THE FERMI-LAT COLLABORATION), arXiv:1303.2733 [astro-ph.HE].
- [30] DUVERNOIS M. A. *et al.* (HEAT COLLABORATION), *Astrophys. J.*, **559** (2001) 296.
- [31] AGUILAR M. *et al.* (AMS COLLABORATION), *Phys. Rep.*, **366** (2002) 331.
- [32] GRASSO D., PROFUMO S., STRONG A. W., BALDINI L., BELLAZZINI R., BLOOM E. D., BREGEON J., DI BERNARDO G., GAGGERO D., GIGLIETTO N., KAMAE T., LATRONICO L., LONGO F., MAZZIOTTA M. N., MOISEEV A. A., MORSELLI A., ORMES J. F., PESCE-ROLLINS M., POHL M., RAZZANO M., SGRO C., SPANDRE G. and STEPHENS T. E., *Astropart. Phys.*, **32** (2009) 140 arXiv:0905.0636.
- [33] ACKERMANN M. *et al.* (FERMI COLLABORATION), *Phys. Rev. D*, **82** (2010) 092003 arXiv:1008.5119.
- [34] AGUILAR M. *et al.* (AMS-02 COLLABORATION), *Phys. Rev. Lett.*, **110** (2013) 141102.