

Combined CP violation measurements by the BaBar and Belle Collaborations

M. RÖHRKEN on behalf of the BABAR and BELLE COLLABORATIONS
California Institute of Technology - Pasadena, CA, USA

received 26 July 2016

Summary. — We present a measurement of the time-dependent CP violation of $\bar{B}^0 \rightarrow D_{CP}^{(*)} h^0$ decays, where the light neutral hadron h^0 is a π^0 , η or ω meson, and the neutral D meson decays to the two-body CP eigenstates $K^+ K^-$, $K_S^0 \pi^0$ or $K_S^0 \omega$. The measurement is performed by combining the final data sets of $(471 \pm 3) \times 10^6$ $B\bar{B}$ pairs collected by the BABAR experiment and $(772 \pm 11) \times 10^6$ $B\bar{B}$ pairs by the Belle experiment in a single physics analysis. In this first measurement performed on a data sample of more than 1 ab^{-1} , we report a first observation of CP violation in $\bar{B}^0 \rightarrow D_{CP}^{(*)} h^0$ decays governed by mixing-induced CP violation according to the weak phase β . We measure the CP asymmetry parameters $-\eta_f \mathcal{S} = +0.66 \pm 0.10$ (stat.) ± 0.06 (syst.) and $\mathcal{C} = -0.02 \pm 0.07$ (stat.) ± 0.03 (syst.).

In the Standard Model, the only source of CP violation is an irreducible complex phase in the three-family Cabibbo-Kobayashi-Maskawa (CKM) quark-mixing matrix [1, 2]. The B factory experiments BABAR at SLAC in the US and Belle at KEK in Japan established CP violation in the neutral and charged B meson system [3-6]. Both experiments precisely measured the angle β of the Unitarity Triangle, defined by the CKM matrix elements V_{ij} as $\arg[-V_{cd}V_{cb}^*/V_{td}V_{tb}^*]$, using the time-dependent CP violation in $b \rightarrow c\bar{c}s$ transitions [7, 8]. The current world average is $\sin(2\beta) = 0.68 \pm 0.02$ [9]. The time-dependent CP violation of $b \rightarrow c\bar{c}s$ transitions is associated with theoretical uncertainties due to possible penguin contributions.

A complementary and theoretically clean probe for β is provided by $\bar{B}^0 \rightarrow D^{(*)} h^0$ decays, where $h^0 \in \{\pi^0, \eta, \omega\}$ denotes a light neutral hadron. These decays are mediated only by tree-level amplitudes that are dominated by CKM-favored $b \rightarrow c\bar{u}d$ transitions. In $\bar{B}^0 \rightarrow D^{(*)} h^0$ decays, an interference between the decay amplitudes with and without B^0 - \bar{B}^0 oscillations emerges if the neutral D meson decays to a CP eigenstate D_{CP} . In this case, the time evolution of $\bar{B}^0 \rightarrow D_{CP}^{(*)} h^0$ decays is governed only by the weak phase β [10]. The measurement of the time-dependent CP violation of $\bar{B}^0 \rightarrow D_{CP}^{(*)} h^0$ decays allows for a determination of β that is theoretically more clean than using $b \rightarrow c\bar{c}s$ transitions, and can provide a new gold standard reference for the new physics searches

in the mixing-induced CP violation of $b \rightarrow s$ penguin-mediated B meson decays. Any sizable deviation in the CP violation of $\bar{B}^0 \rightarrow D_{CP}^{(*)}h^0$ decays from $b \rightarrow s$ penguin transitions would point to physics beyond the Standard Model which could be caused, for example, by unobserved heavy particles or additional CP violating phases contributing to loop diagrams in $b \rightarrow s$ penguin transitions [11].

However, the measurements of $\bar{B}^0 \rightarrow D^{(*)}h^0$ decays are experimentally challenging because of low B and D meson branching fractions ($\mathcal{O}(10^{-4})$ and $\mathcal{O}(\leq 10^{-2})$, respectively), low reconstruction efficiencies, and large backgrounds. Previous measurements carried out separately by the BABAR and Belle collaborations using two-body [12] and three-body D meson decays [13, 14] were not sensitive enough to establish CP violation in $\bar{B}^0 \rightarrow D^{(*)}h^0$ decays.

In this article, we present a time-dependent CP violation measurement of $\bar{B}^0 \rightarrow D_{CP}^{(*)}h^0$ decays. In this measurement, we combine for the first time the large final data sets of $(471 \pm 3) \times 10^6 B\bar{B}$ pairs collected by the BABAR experiment and $(772 \pm 11) \times 10^6 B\bar{B}$ pairs by the Belle experiment in a single physics analysis. The novel approach increases the achievable experimental precision and enables time-dependent CP violation measurements in the neutral B meson system with unprecedented sensitivity.

In $\bar{B}^0 \rightarrow D_{CP}^{(*)}h^0$ decays, the light neutral hadron h^0 is reconstructed as a π^0 , η or ω meson and the neutral D meson is reconstructed in the decays to two-body CP eigenstates K^+K^- , $K_S^0\pi^0$ and $K_S^0\omega$. D^{*0} mesons are reconstructed in the decays to $D^0\pi^0$. In the measurement, twelve final states are reconstructed in total, among them seven (five) CP -even (CP -odd) states. Continuum $e^+e^- \rightarrow q\bar{q}$ ($q \in \{u, d, s, c\}$) events are the dominant source of background. This background is reduced by selection requirements on multivariate classifiers provided by neural networks trained on variables characterizing the event shapes. The $\bar{B}^0 \rightarrow D_{CP}^{(*)}h^0$ signal yields are estimated by unbinned maximum likelihood fits to the beam-constrained mass defined as $M_{bc} \equiv \sqrt{(E_{\text{beam}}^*/c^2)^2 - (p_B^*/c)^2}$, where E_{beam}^* is the energy of the beam and p_B^* is the momentum of reconstructed B meson candidates in the e^+e^- center-of-mass frame. A signal yield of 508 ± 31 (757 ± 44) events is obtained for BaBar (Belle). The signal yields split into decay modes and experiments are summarized in table I. The M_{bc} data distributions and projections of the fits are shown in fig. 1.

The time-dependent CP violation analysis is performed by combining the BaBar and Belle flavor-tagged proper decay time distributions on the likelihood level. In the

TABLE I. – Summary of $\bar{B}^0 \rightarrow D_{CP}^{(*)}h^0$ signal yields.

Decay mode	BaBar	Belle
$\bar{B}^0 \rightarrow D_{CP}\pi^0$	241 ± 22	345 ± 25
$\bar{B}^0 \rightarrow D_{CP}\eta$	106 ± 14	148 ± 18
$\bar{B}^0 \rightarrow D_{CP}\omega$	66 ± 10	151 ± 17
$\bar{B}^0 \rightarrow D_{CP}^*\pi^0$	72 ± 12	80 ± 14
$\bar{B}^0 \rightarrow D_{CP}^*\eta$	39 ± 8	39 ± 10
$\bar{B}^0 \rightarrow D_{CP}^{(*)}h^0$ total	508 ± 31	757 ± 44




Fig. 1. – The M_{bc} data distributions (data points with error bars) and projections of the fits (solid line) of $\bar{B}^0 \rightarrow D_{CP}^{(*)} h^0$ decays for (a) BaBar and (b) Belle. The dashed and dotted lines represent projections of the signal and background components of the fits, respectively.

measurement, the following log-likelihood function is maximized:

$$(1) \quad \ln \mathcal{L} = \sum_i \ln \mathcal{P}_i^{BaBar} + \sum_j \ln \mathcal{P}_j^{Belle}.$$

The indices i and j denote events reconstructed from BaBar and Belle data, respectively. The probability density functions for signal are constructed from the convolution of experiment-specific resolution functions accounting for the finite B meson vertex resolution with the decay rate of a neutral B meson decaying to a CP eigenstate, defined as

$$(2) \quad g(\Delta t) = \frac{e^{-|\Delta t|/\tau_{B^0}}}{4\tau_{B^0}} \{1 + q[S \sin(\Delta m_d \Delta t) - \mathcal{C} \cos(\Delta m_d \Delta t)]\},$$

where Δt is the proper time interval between the decays of the two B mesons produced in an $\Upsilon(4S)$ decay, and S and \mathcal{C} , respectively, measure mixing-induced and direct CP violation. The $q = +1$ (-1) denotes the b -flavor content when the accompanying B meson is tagged as a B^0 (\bar{B}^0) meson. The neutral B meson lifetime is represented by τ_{B^0} , and the $B^0 - \bar{B}^0$ mixing frequency by Δm_d . The experimental flavor-tagged proper decay time distributions for BaBar and Belle for the CP -even and CP -odd final states and projections of the fit are shown in fig. 2.

The result of the measurement including statistical and systematic uncertainties is

$$(3) \quad \begin{aligned} -\eta_{f_{CP}} \mathcal{S} &= +0.66 \pm 0.10 \text{ (stat.)} \pm 0.06 \text{ (syst.)} \\ \mathcal{C} &= -0.02 \pm 0.07 \text{ (stat.)} \pm 0.03 \text{ (syst.)}, \end{aligned}$$

where $\eta_{f_{CP}}$ is the CP eigenvalue of the final state. The individual sources contributing to the systematic uncertainties are summarized in table II.




Fig. 2. – The proper time interval distributions of $\bar{B}^0 \rightarrow D_{CP}^{(*)} h^0$ decays (data points with error bars) for B^0 tags (red) and \bar{B}^0 tags (blue) and the corresponding CP violating asymmetries for (a)-(b) BaBar and (c)-(d) Belle for candidates associated with high-quality flavor tags.

TABLE II. – *Systematic uncertainties of the measured time-dependent CP violation parameters of $\bar{B}^0 \rightarrow D_{CP}^{(*)} h^0$ decays (in units of 10^{-2}).*

Source	S	C
Background Δt PDFs	0.4	0.1
Δt resolution functions	2.0	0.4
Flavor-tagging	0.3	0.3
Peaking background	4.9	0.9
Physics parameters	0.2	< 0.1
Possible fit bias	0.6	0.8
Signal purity	0.6	0.3
Tag-side interference	0.1	1.4
Vertex reconstruction	1.5	1.4
Total	5.6	2.5

The statistical significance of the result is estimated by a likelihood-ratio approach. Including systematic uncertainties, the measurement excludes the hypothesis of no mixing-induced CP violation in $\bar{B}^0 \rightarrow D_{CP}^{(*)}h^0$ decays at a confidence level of $1 - 6.6 \times 10^{-8}$. This corresponds to a significance of 5.4 standard deviations, establishing an observation of CP violation in $\bar{B}^0 \rightarrow D_{CP}^{(*)}h^0$ decays for the first time. The result agrees well with the Standard Model prediction of $\mathcal{S} = -\eta_{f_{CP}} \sin(2\beta)$ and $\mathcal{C} = 0$. The measured mixing-induced CP violation is compatible with the current world average of $\sin(2\beta) = 0.68 \pm 0.02$ [9] within 0.2 standard deviations.

In summary, we combine for the first time the final data sets of the BaBar and Belle experiments in a single physics analysis to measure the time-dependent CP violation of $\bar{B}^0 \rightarrow D_{CP}^{(*)}h^0$ decays. In this first measurement performed on a data sample of more than 1 ab^{-1} , we observe for the first time CP violation in $\bar{B}^0 \rightarrow D_{CP}^{(*)}h^0$ decays driven by mixing-induced CP violation, and we measure $\sin(2\beta) = 0.66 \pm 0.10$ (stat.) ± 0.06 (syst.). The result is consistent within 0.2 standard deviations with the more precise world average of $\sin(2\beta) = 0.68 \pm 0.02$ [9] measured from $b \rightarrow c\bar{c}s$ transitions. At the expected precision of the upcoming high-luminosity B factory experiment Belle II, $\bar{B}^0 \rightarrow D_{CP}^{(*)}h^0$ decays will provide a new new gold standard reference for the new physics searches in the mixing-induced CP violation of $b \rightarrow s$ penguin-mediated B meson decays.

REFERENCES

- [1] CABIBBO N., *Phys. Rev. Lett.*, **10** (1963) 531.
- [2] KOBAYASHI M. and MASKAWA T., *Prog. Theor. Phys.*, **49** (1973) 652.
- [3] BABAR COLLABORATION (AUBERT B. *et al.*), *Phys. Rev. Lett.*, **87** (2001) 091801.
- [4] BELLE COLLABORATION (ABE K. *et al.*), *Phys. Rev. Lett.*, **87** (2001) 091802.
- [5] BABAR COLLABORATION (AUBERT B. *et al.*), *Phys. Rev. Lett.*, **93** (2004) 131801.
- [6] BELLE COLLABORATION (CHAO Y. *et al.*), *Phys. Rev. Lett.*, **93** (2004) 191802.
- [7] BABAR COLLABORATION (AUBERT B. *et al.*), *Phys. Rev. D*, **79** (2009) 072009.
- [8] BELLE COLLABORATION (ADACHI I. *et al.*), *Phys. Rev. Lett.*, **108** (2012) 171802.
- [9] HEAVY FLAVOR AVERAGING GROUP (AMHIS Y. *et al.*), arXiv:1412.7515 (2014).
- [10] FLEISCHER R., *Phys. Lett. B*, **562** (2003) 234; FLEISCHER R., *Nucl. Phys. B*, **659** (2003) 321.
- [11] GROSSMAN Y. and WORAH M. P., *Phys. Lett. B*, **395** (1997) 241; LONDON D. and SONI A., *Phys. Lett. B*, **407** (1997) 61; MOROI T., *Phys. Lett. B*, **493** (2000) 366; BAEK S., GOTO T., OKADA Y. and OKUMURA K. I., *Phys. Rev. D*, **64** (2001) 095001; CHANG D., MASIERO A. and MURAYAMA H., *Phys. Rev. D*, **67** (2003) 075013.
- [12] BABAR COLLABORATION (AUBERT B. *et al.*), *Phys. Rev. Lett.*, **99** (2007) 081801.
- [13] BELLE COLLABORATION (KROKOVNY P. *et al.*), *Phys. Rev. Lett.*, **97** (2006) 081801.
- [14] BABAR COLLABORATION (AUBERT B. *et al.*), *Phys. Rev. Lett.*, **99** (2007) 231802.