

# Spatial distribution of *Biomphalaria* spp., the intermediate host snails of *Schistosoma mansoni*, in Brazil

Ronaldo G.C. Scholte<sup>1,2,3</sup>, Omar S. Carvalho<sup>3</sup>, John B. Malone<sup>4</sup>, Jürg Utzinger<sup>1,2</sup>, Penelope Vounatsou<sup>1,2</sup>

<sup>1</sup>Department of Epidemiology and Public Health, Swiss Tropical and Public Health Institute, P.O. Box, CH-4002 Basel, Switzerland; <sup>2</sup>University of Basel, P.O. Box, CH-4003 Basel, Switzerland; <sup>3</sup>Centro de Pesquisas René Rachou/Fiocruz-MG, Av Augusto de Lima 1715, Barro Preto, Belo Horizonte, CEP:30190-002, MG, Brazil; <sup>4</sup>Department of Pathobiological Sciences, School of Veterinary Medicine, Louisiana State University, Baton Rouge, LA 70803, USA

**Abstract.** Schistosomiasis mansoni remains an important parasitic disease of man, endemic in large parts of sub-Saharan Africa, the Middle East, South America and the Caribbean. The aetiological agent is the trematode *Schistosoma mansoni*, whereas aquatic snails of the genus *Biomphalaria* act as intermediate hosts in the parasite life cycle. In Brazil, the distribution of *Biomphalaria* spp. is closely associated with the occurrence of schistosomiasis. The purpose of this study was to map and predict the spatial distribution of the intermediate host snails of *S. mansoni* across Brazil. We assembled snail “presence-only” data and used a maximum entropy approach, along with climatic and environmental variables to produce predictive risk maps. We identified a series of risk factors that govern the distribution of *Biomphalaria* snails. We find that high-risk areas for *B. glabrata* are concentrated in the regions of Northeast and Southeast and the northern part of the South region. *B. straminea* are found in the Northeast and Southeast regions, and *B. tenagophila* are concentrated in the Southeast and South regions. Our findings confirm that the presence of the intermediate host snails is correlated with the occurrence of schistosomiasis mansoni. The generated risk maps of intermediate host snails might assist the national control programme for spatial targeting of control interventions and to ultimately move towards schistosomiasis elimination in Brazil.

**Keywords:** *Biomphalaria*, intermediate host snail, schistosomiasis, risk mapping and prediction, ecological niche model, maximum entropy, Brazil.

## Introduction

Schistosomiasis mansoni is a human parasitic disease, currently endemic in 54 countries in Africa, the Middle East, South America and the Caribbean (Chitsulo et al., 2000; Gryseels et al. 2006; Steinmann et al., 2006). In Brazil, Katz and Peixoto (2000), using data from the National Health Foundation, estimated that 6.4 million people in 18 states are affected by schistosomiasis. The causative agent is the blood-dwelling fluke *Schistosoma mansoni*, and the intermediate hosts are aquatic snails of the genus *Biomphalaria* (Mollusca: Gastropoda: Pulmonata: Planorbidae). Thus far, 11 species and one subspecies of *Biomphalaria* have been described in Brazil. Among these, three species (i.e. *B. glabrata*, *B. tenagophila*

and *B. straminea*) were found naturally infected with *S. mansoni*. Another three species (i.e. *B. amazonica*, *B. peregrina* and *B. cousini*) have been considered as potential intermediate hosts (Corrêa and Paraense, 1971; Paraense and Corrêa, 1973; Caldeira et al., 2010, Teodoro et al., 2010).

Clearly, *B. glabrata* is recognised as the best adapted intermediate host for transmitting *S. mansoni* due to its wide geographical distribution, high rates of infection and transmission efficiency. The distribution of this snail species is almost always associated with the occurrence of schistosomiasis mansoni, the observation of this fact was first mentioned by Lutz (1917) (Paraense, 1983; Teles and Vaz, 1987, Carvalho et al., 2008). It follows that maps of the distribution of *B. glabrata* and other (potential) intermediate hosts could serve as an important tool for spatial targeting of schistosomiasis control interventions (Stensgaard et al., 2012). However, only few studies assessed the distribution of *Biomphalaria* spp. in Brazil (Malone et al., 2005; Guimarães et al., 2009; Baboza et al., 2012). It should be noted that available data on *Biomphalaria* spp. in Brazil is “presence-only”, while care is indicat-

Corresponding author:  
Ronaldo G.C. Scholte  
Department of Epidemiology and Public Health  
Swiss Tropical and Public Health Institute  
P.O. Box, CH-4002 Basel, Switzerland  
Tel. +41 61 284 8226; Fax +41 61 284 8105  
E-mail: ronaldo.scholte@unibas.ch

ed in absence data, since surveyed areas are large and the surveyors might have missed sites where snails occurred but were not discovered due to the limited sampling effort.

When both presence and absence data are available, classical statistical approaches can be used to analyse the data (Corsi et al., 2000; Guisan and Zimmerman, 2000; Elith, 2002; Scott et al., 2002). However, for presence-only data, modelling techniques are required that take into account the lack of zeros in the data in order to obtain accurate estimates of the outcome distribution. Different methods have been used for presence-only data when modelling species distribution, such as ecological niche models (ENMs). Among others, prominent ENMs include generalized linear models (GLMs), generalized additive models (GAMs), generalized linear mixed models (GLMMs) (Fenton et al., 2010), BIOCLIM (Busby, 1986; Nix, 1986), DOMAIN (Carpenter et al., 1993), environmental niche factor analysis (ENFA) (Hirzel et al., 2002), genetic algorithm for rule-set prediction (GARP) (Stockwell and Noble, 1992) and maximum entropy (MaxEnt) models (Phillips et al. 2004). MaxEnt models proved particularly useful, as they pursue a general purpose method for making predictions from presence-only data (Ponder et al., 2001; Anderson et al., 2002, 2003; Graham et al., 2004; Philips et al., 2006; Stensgaard et al., 2012).

In this study, we employed compiled *Biomphalaria* spp. presence-only data from Brazil to produce predictive risk maps of the spatial distribution of the intermediate host snails of *S. mansoni*. We used a MaxEnt modelling approach using as predictors both environmental and climatic proxies readily obtained from remote sensing and climate databases. The maps presented here will be useful for the spatial targeting of schistosomiasis control interventions and future efforts emphasising schistosomiasis elimination.

## Materials and methods

### Malacological data

We obtained *Biomphalaria* spp. occurrence data (i.e. presence-only) for Brazil from two main sources. First, data were readily available from the Laboratory of Helminthiasis and Medical Malacology (LHMM) of the René Rachou Research Center (CPqRR/Fiocruz-MG). Of note, the LHMM is a reference laboratory for schistosomiasis in Brazil, performing the examination and identification of the snails of the genus *Biomphalaria* according to the demands of health services. Second, data were retrieved through a systematic review of the peer-reviewed literature and from grey literature (Carvalho et al., 2008). Our database comprises 1,977 municipalities, hence 35.5% from all the municipalities in Brazil (Fig. 1).

### Environmental and climatic data

Environmental and climatic proxies were considered in our analyses, since these are the main predictors for the distribution of *Biomphalaria* spp. Environmental data were extracted from different freely accessible remote sensing data sources, as summarised in Table 1. In brief, land surface temperature (LST) data were used as a proxy for day and night temperature. The normalized difference vegetation index (NDVI) was employed as a proxy for moisture. A digital elevation model (DEM) was used to extract altitude data.

Climatic data were obtained from Worldclim-Global Climate Data, which provides interpolated monthly climatic information from weather stations averaged over a 50-year period (from 1950 to 2000) at a spatial resolution of 1 km (Table 1). The Worldclim data have been used successfully in ENMs and by the global climate change community (Waltari et al., 2007).




Fig. 1. Observed spatial distribution of the intermediate host snails of *S. mansoni* in Brazil; (A) *B. glabrata*; (B) *B. straminea*; (C) *B. tenagophila* and (D) *Biomphalaria* spp. (Carvalho et al., 2008).

Table 1. Data sources and properties of the climatic and other environmental covariates used in our models to predict the occurrence of *Biomphalaria* spp. in Brazil.

Source	Data type	Data period	Temporal resolution	Spatial resolution
Shuttle Radar Topography Mission (SRTM) data	Digital elevation model (DEM)	2000	Once	1 km
Moderate Resolution Imaging Spectroradiometer (MODIS)/Terra	Land surface temperature (LST) for day and night	2005-2009	8 days	1 km
	Normalized difference vegetation index (NDVI)	2005-2009	16 days	1 km
	BIO1 (annual mean temperature)	1950-2000	Once	1 km
	BIO2 (mean diurnal range (mean of monthly (max temp - min temp)))	1950-2000	Once	1 km
	BIO3 (isothermality (P2/P7) (* 100))	1950-2000	Once	1 km
	BIO4 (temperature seasonality (standard deviation *100))	1950-2000	Once	1 km
	BIO5 (max. temperature of warmest month)	1950-2000	Once	1 km
	BIO6 (min. temperature of coldest month)	1950-2000	Once	1 km
	BIO7 (temperature annual range (P5-P6))	1950-2000	Once	1 km
	BIO8 (mean temperature of wettest quarter)	1950-2000	Once	1 km
	BIO9 (mean temperature of driest quarter)	1950-2000	Once	1 km
Worldclim Global Climate	BIO10 (mean temperature of warmest quarter)	1950-2000	Once	1 km
	BIO11 (mean temperature of coldest quarter)	1950-2000	Once	1 km
	BIO12 (annual precipitation)	1950-2000	Once	1 km
	BIO13 (precipitation of wettest month)	1950-2000	Once	1 km
	BIO14 (precipitation of driest month)	1950-2000	Once	1 km
	BIO15 (precipitation seasonality (coefficient of variation))	1950-2000	Once	1 km
	BIO16 (precipitation of wettest quarter)	1950-2000	Once	1 km
	BIO17 (precipitation of driest quarter)	1950-2000	Once	1 km
	BIO18 (precipitation of warmest quarter)	1950-2000	Once	1 km
	BIO19 (precipitation of coldest quarter)	1950-2000	Once	1 km

### MaxEnt modelling approach

We used a MaxEnt modelling approach proposed by Phillips et al. (2006) for prediction of the intermediate host snail distribution. The MaxEnt is an iterative method used to obtain predictions or make inferences from incomplete information (e.g. presence-only data). The main purpose is to estimate the unknown probability distribution of a species based on the principle of maximum entropy (Phillips et al., 2006). Entropy, as defined by Shannon (1948), is “a measure of how much “choice” is involved in the selection of an event”. A distribution with higher entropy, involves more choices. Given a set of samples (e.g. presence of species) and a set of features (e.g. environmental/climatic variables), the MaxEnt model estimates niches by finding the distribution of probabilities closest to uniform (maximum entropy), constrained by the fact that feature values match their empirical average. The final result is a model that predicts areas where the snails are most likely to occur based on environmental and/or climatic correlations.

We employed the MaxEnt method in the present study to generate probability surfaces of the geographical distributions of the (potential) intermediate

host snails of *S. mansoni*. The method’s principle is to estimate the spatial distribution of the snails of the genus *Biomphalaria* that are most probable to be present in a certain area, constrained to known observations (i.e. presence of intermediate hosts of *S. mansoni*). MaxEnt uses entropy as the means to generalize specific observations pertaining to the presence of a particular species.

The area under the curve (AUC) test statistics was used to measure the accuracy of the predictive distribution models. It estimates the probability of a test pixel to be correctly predicted as suitable for species presence as opposed to the probability of a randomly selected pixel of the map. The AUC ranges from zero to one, where values near one indicate high predictive performance of the model, while values smaller than 0.5 indicate low model predictive ability (Wiley et al., 2003; Allouche et al., 2006; Elith et al., 2006).

To determine the spatial distribution of the intermediate host snails of *S. mansoni* in Brazil, four models were constructed. The models were: (i) *B. glabrata*, (ii) *B. straminea*, (iii) *B. tenagophila*, and (iv) *Biomphalaria* spp., the latter including all three intermediate host species.

### Statistical analysis

We used the MaxEnt software version 3.3.3e (Princeton University; Princeton, NJ, USA) to perform our analysis. Visualization and geostatistical display was carried out in ArcGIS version 9.3 (ESRI; Redlands, CA, USA). The final models were plotted with a 1 km spatial resolution.

### Results

Fig. 2A shows the most suitable areas for the presence of *B. glabrata* across Brazil. The following environmental and climatic variables showed the highest contribution to the predictive model: temperature seasonality, maximum temperature of warmest month, annual precipitation and monthly mean diurnal range temperature. Our model shows a high probability of

the occurrence of *B. glabrata* in the regions of Northeast and Southeast and the northern part of the South regions within Brazil.

The most suitable areas for the presence of *B. straminea* are shown in Fig. 2B. The map indicates high probability of occurrence of *B. straminea* in the Northeast and Southeast regions of Brazil. Annual precipitation, temperature seasonality and temperature annual range were the three variables with the highest contribution to the predictive ability of the model.

The prediction map in Fig. 2C shows that, the areas which are likely for *B. tenagophila* to be present, are concentrated in the Southeast and South regions of Brazil. Temperature seasonality, isothermality and monthly mean temperature of driest quarter were the most important predictors of the species' geographical distribution.




Fig. 2. Probability of the spatial distribution of the intermediate host snails of *S. mansoni* in Brazil; (A) *B. glabrata*; (B) *B. straminea*; (C) *B. tenagophila* and (D) *Biomphalaria* spp.



Model-based predictions suggest that the areas most suitable for the presence of *Biomphalaria* spp. are located in the Northeast, Southeast and South regions of Brazil (Fig. 2D). The climatic parameters related to the species distribution are temperature seasonality, annual precipitation and monthly mean diurnal range.

The predictive ability of the above models were high with an AUC equal to 0.836, 0.859, 0.863 and 0.901 for *Biomphalaria* spp., *B. straminea*, *B. tenagophila* and *B. glabrata*, respectively.

## Discussion

To our knowledge, we present the first predictive maps, using MaxEnt, of the spatial distribution of *B. glabrata*, the key intermediate host snail of *S. mansoni*, in Brazil. This snail species shows a wide distribution with a high probability of presence in the Northeast and Southeast regions and the northern part of the South region. Moreover, we present spatially explicit probability maps of the presence of *B. tenagophila* and *B. straminea*, which have also been identified as intermediate host snails of *S. mansoni* in Brazil. While *B. straminea* is concentrated in the Northeast region, *B. tenagophila* primarily occurs in the Southeast and South regions. A climatic suitability map showed that the region extending from Northeast to South has high probability of snail occurrence irrespective of their species with the highest one over 50% estimated in the Northeast, Southeast and part of South regions of Brazil. Our results agree with the literature showing the same pattern for each snail species (Carvalho et al., 2008; Barboza et al., 2012).

We employed a MaxEnt modelling approach, which has been proposed as a particularly useful ENM, to analyse a comprehensive dataset with more than 1,900 presence-only data points for *Biomphalaria* (used as outcome measure) in relation to climatic and environmental factors (used as covariates). ENMs have been widely used to address questions of species distribution and ecology in biodiversity-related studies using presence-only data (Kadmon and Heller, 1998; Gottfried et al., 1999; Peterson et al., 1999, 2001, 2002; Bakkenes et al., 2002; Peterson and Shaw, 2003; Phillips et al., 2006). More recently, ENMs, and specifically MaxEnt models, have been applied to deepen our understanding of vector and intermediate host distribution (Costa et al., 2002; Batista and Gurgel-Gonçalves, 2009; Stensgaard et al., 2012). However, only few studies used ENMs to predict the distribution of the intermediate host snails of schistosomiasis, with a notable exception of recent predictive

risk maps of *Biomphalaria* spp. in Africa (Stensgaard et al., 2012).

In all our models, both temperature and precipitation were identified as important environmental features governing the distribution of *Biomphalaria* spp. in Brazil. Indeed, our models showed high predictive accuracy with AUC values above 0.8. According to Phillips and Dudik (2008), AUC values above 0.75 can be considered as to fall within the best model-fit category.

Noteworthy issues arising from our work can be summarised as follows. First, our results confirm that the distribution of aquatic snails is governed by environmental and climatic features, and hence the Worldclim database is useful for ENMs. Second, the MaxEnt model employed in this study proved useful to analyse and predict presence-only data for understanding geographical and ecological distributions of species in disease transmission cycles. Third, our maps of probability of the intermediate host snails' distribution follow the same pattern of the predicted prevalence of schistosomiasis in Brazil. Hence, the *Biomphalaria* distribution map developed here might be used as predictor for *S. mansoni* distribution. Since the presence of the intermediate host snails of *S. mansoni* is a proxy for the disease distribution, and it is widely acknowledged which species is the most adapted intermediate host, the methodology and the generated risk maps are useful for decision makers to delineate priority areas for schistosomiasis control interventions, so that limited resources can be allocated most effectively. Our maps can guide the spatial targeting of control interventions and are of particular relevance now that efforts are getting under way to eliminate schistosomiasis in Brazil and elsewhere (Rollinson et al., 2012).

Some important issues related to the nature and precision of the *Biomphalaria* species data need to be considered when interpreting our findings. The *Biomphalaria* species data were extracted from peer-reviewed articles and grey literature. In most cases the data records were not georeferenced, and hence no precise spatial information is available. Most of the time, the data were at municipality level. Our assumption then was that the species found in a given municipality is uniformly distributed within this municipality along the drainage network. It would be interesting to conduct malacological surveys in selected municipalities, and georeference the occurrence of snails in order to determine whether data at higher spatial resolution would further improve model accuracy. Since a global neglected tropical disease (GNTD) database is now

available (Hürlimann et al., 2011) and spatially explicit parasite prevalence data (e.g. *S. mansoni*) can be freely obtained, it will be important to further populate the GNTD database with information on occurrence and density of intermediate host snails (and vectors for other diseases) to assist global control and elimination efforts targeting schistosomiasis and other neglected tropical diseases.

### Acknowledgements

This investigation received financial support from the UBS Optimus Foundation, the Swiss-Brazilian joint research programme (BJRP 011008) and CNPq (Process 150386/2012-5).

### References

- Allouche O, Tsoar A, Kadmon R, 2006. Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). *J Appl Ecol* 43, 1223-1232.
- Anderson RP, Gomez-Laverde M, Peterson AT, 2002. Geographical distributions of spiny pocket mice in South America: insights from predictive models. *Global Ecol Biogeogr* 11, 131-141.
- Anderson RP, Lew D, Peterson AT, 2003. Evaluating predictive models of species' distributions: criteria for selecting optimal models. *Ecol Modell* 162, 211-232.
- Bakkenes M, Alkemade JRM, Ihle F, Leemansand R, Latour JB, 2002. Assessing effects of forecasted climate change on the diversity and distribution of European higher plants for 2050. *Glob Change Biol* 8, 390-407.
- Barboza DM, Rollemberg CVV, de Amorim FJR, Zhang C, de Melo CM, de Lurdes Sierpe Jeraldo V, Santos NC, Silva MMBL, de Almeida JAP, Ueta MT, de Jesus AR, 2012. *Biomphalaria* species distribution and its effect on human *Schistosoma mansoni* infection in an irrigated area used for rice cultivation in northeast Brazil. *Geospat Health* 6, S103-S109.
- Batista TA, Gurgel-Gonçalves R, 2009. Ecological niche modeling and differentiation between *Rhodnius neglectus* Lent, 1954 and *Rhodnius nasutus* Stål, 1859 (Hemiptera: Reduviidae: Triatominae) in Brazil. *Mem Inst Oswaldo Cruz* 104, 1165-1170.
- Busby JR, 1986. A biogeographical analysis of *Nothofagus cunninghamii* (Hook.) Oerst. in southeastern Australia. *Aust J Ecol* 11, 1-7.
- Caldeira RL, Teodoro TM, Gomes MFB, Carvalho OS, 2010. Preliminary studies investigating the occurrence of *Biomphalaria cousini* in Brazil. *Mem Inst Oswaldo Cruz* 105, 485-487.
- Carpenter G, Gillison AN, Winter J, 1993. DOMAIN: a flexible modeling procedure for mapping potential distributions of plants and animals. *Biodivers Conserv* 2, 667-680.
- Carvalho OS, Amaral RS, Dutra LV, Scholte RGC, Guerra M, 2008. Distribuição espacial de *Biomphalaria glabrata*, *B. straminea* e *B. tenagophila* moluscos hospedeiros intermediários do *Schistosoma mansoni* no Brasil. In: Carvalho OS, Coelho PMZ, Lenzi HL (eds). *Schistosoma mansoni* e esquistossomose: uma visão multidisciplinar. Rio de Janeiro: Fiocruz, v. 1, pp. 395-418.
- Chitsulo L, Engels D, Montresor A, Savioli L, 2000. The global status of schistosomiasis and its control. *Acta Trop* 77, 41-51.
- Corrêa LR, Paraense WL, 1971. Susceptibility of *Biomphalaria amazonica* to infection with two strains of *Schistosoma mansoni*. *Rev Inst Med Trop São Paulo*, 13, 387-390.
- Corsi F, de Leeuw J, Skidmore A, 2000. Modeling species distribution with GIS. In: Boitani L, Fuller T (eds). New York: Columbia University Press, pp. 389-434.
- Costa J, Peterson AT, Beard CB, 2002. Ecological niche modeling and differentiation of populations of *Triatoma brasiliensis* Neiva, 1911, the most important Chagas disease vector in north-eastern Brazil (Hemipter, Reduviidae, Triatominae). *Am J Trop Med Hyg* 67, 516-520.
- Elith J, 2002. Quantitative methods for modeling species habitat: comparative performance and an application to Australian plants. In: Ferson S, Burgman M (eds). *Quantitative methods for conservation biology*. Springer-Verlag, New York, pp. 39-58.
- Elith J, Graham CH, Anderson RP, Dudik M, Ferrier S, Guisan A, Hijmans RJ, Huettmann F, Leathwick JR, Lehmann A, Li J, Lohmann LG, Loiselle BA, Manion G, Moritz C, Nakamura M, Nakazawa Y, Overton JMcC, Peterson AT, Phillips SJ, Richardson KS, Scachetti-Pereira R, Schapire RE, Soberón J, Williams S, Wisz MS, Zimmermann NE, 2006. Novel methods improve prediction of species' distributions from occurrence data. *Ecography* 29, 129-151.
- Fenton A, Viney ME, Lello J, 2010. Detecting interspecific macroparasite interactions from ecological data: patterns and process. *Ecol Lett* 13, 606-615.
- Gottfried M, Pauli H, Reiter K, Grabherr G, 1999. A fine-scaled predictive model for changes in species distribution patterns of high mountain plants induced by climate warming. *Divers Distrib* 5, 241-251.
- Graham CH, Ferrier S, Huettman F, Moritz C, Peterson AT, 2004. New developments in museum-based informatics and applications in biodiversity analysis. *Trends Ecol Evol* 19, 497-503.
- Gryseels B, Polman K, Clerinx J, Kestens L, 2006. Human schistosomiasis. *Lancet* 368, 1106-1118.
- Guimarães RJPS, Freitas CC, Dutra LV, Felgueiras CA, Moura ACM, Amaral RS, Drummond SC, Scholte RGC, Oliveira G, Carvalho OS, 2009. Spatial distribution of *Biomphalaria* mollusks at Sao Francisco River Basin, Minas Gerais, Brazil, using geostatistical procedures. *Acta Trop* 109, 181-186.
- Guisan A, Zimmermann NE, 2000. Predictive habitat distribu-

- tion models in ecology. *Ecol Modell* 135, 147-186.
- Hirzel AH, Hausser J, Chessel D, Perrin N, 2002. Ecological-niche factor analysis: how to compute habitat-suitability maps without absence data? *Ecology* 83, 2027-2036.
- Hürlimann E, Schur N, Boutsika K, Stensgaard AS, Laserna de Himpsl M, Ziegelbauer K, Laizer N, Camenzind L, Di Pasquale A, Ekpo UF, Simoonga C, Mushinge G, Saarnak CFL, Utzinger J, Kristensen TK, Vounatsou P, 2011. Toward an open-access global database for mapping, control, and surveillance of neglected tropical diseases. *PLoS Negl Trop Dis* 5, e1404.
- Kadmon R, Heller J, 1998. Modelling faunal responses to climatic gradients with GIS: land snails as a case study. *J Biogeogr* 25, 527-539.
- Katz N, Peixoto SV, 2000. Análise crítica da estimativa do número de portadores de esquistossomose mansoni no Brasil. *Rev Soc Bras Med Trop* 33, 303-308.
- Lutz A, 1917. Observações sobre a evolução do *Schistosoma mansoni*. *Revista da Sociedade Brasileira de Ciências* 1, 41-48.
- Malone JB, Bavia ME, Amaral R, McNally KL, Nieto P, Carneiro D, Neves A, 2005. A climate-based prediction of the potential distribution of schistosomiasis in Brazil. In: Confaloneiri UEC, Marino DP (eds). *Use of remote sensing for control of infectious diseases*. SOCGRAF, Rio de Janeiro, Brazil, pp. 91-98.
- Nix HA, 1986. A biogeographic analysis of Australian Elapid snakes. In: Longmore R (ed.). *Atlas of Australian Elapid snakes*. Australian Flora and Fauna Series 8, 4-15.
- Paraense WL, 1983. A survey of planorbid molluscs in the Amazonian region of Brazil. *Mem Inst Oswaldo Cruz* 78, 343-361.
- Paraense WL, Corrêa LR, 1973. Susceptibility of *Biomphalaria peregrina* from Brazil and Ecuador to two strains of *Schistosoma mansoni*. *Rev Inst Med Trop São Paulo* 15, 127-130.
- Peterson AT, Sanchez-Cordero V, Beard CB, Ramsey JM, 2002. Ecologic niche modelling and potential reservoirs for Chagas disease, Mexico. *Emerg Infect Dis* 8, 662-667.
- Peterson AT, Sanchez-Cordero V, Soberón J, Bartley J, Buddemeier RH, Navarro-Siguenza AG, 2001. Effects of global climate change on geographic distributions of Mexican Cracidae. *Ecol Modell* 144, 21-30.
- Peterson AT, Shaw J, 2003. *Lutzomyia* vectors for cutaneous leishmaniasis in southern Brazil: ecological niche models, predicted geographic distributions, and climate change effects. *Int J Parasitol* 33, 919-931.
- Peterson AT, Sober NJ, Sanchez-Cordero V, 1999. Conservatism of ecological niches in evolutionary time. *Science* 285, 1265-1267.
- Phillips SJ, Anderson RP, Schapire RE, 2006. Maximum entropy modeling of species geographic distributions. *Ecol Modell* 190, 231-259.
- Phillips SJ, Dudik M, 2008. Modeling of species distributions with Maxent: new extensions and a comprehensive evaluation. *Ecography* 31, 161-175.
- Phillips SJ, Dudik M, Schapire RE, 2004. A maximum entropy approach to species distribution modeling. In: *Proceedings of the 21<sup>st</sup> International Conference on Machine Learning*, ACM Press, New York, pp. 655-662.
- Ponder WF, Carter GA, Flemons P, Chapman RR, 2001. Evaluation of museum collection data for use in biodiversity assessment. *Conserv Biol* 15, 648-657.
- Rollinson D, Knopp S, Levitz S, Stothard JR, Tchuem Tchuente LA, Garba A, Mohammed KA, Schur N, Person B, Colley DG, Utzinger J, 2012. Time to set the agenda for schistosomiasis elimination. *Acta Trop* (in press; doi:10.1016/j.actatropica.2012.04.013).
- Scott JM, Heglund PJ, Morrison ML, Haufler JB, Raphael MG, Wall WA, Samson FB, 2002. *Predicting species occurrences: issues of accuracy and scale*. Washington, DC: Island Press.
- Shanon C, 1948. A mathematical theory of communication. *Bell Syst Tech J* 27, 379-423, 623-656.
- Steinmann P, Keiser J, Bos R, Tanner M, Utzinger J, 2006. Schistosomiasis and water resources development: systematic review, meta-analysis, and estimates of people at risk. *Lancet Infect Dis* 6, 411-425.
- Stensgaard AS, Utzinger J, Vounatsou P, Hürlimann E, Schur N, Saarnak CFL, Mushinge G, Simoonga C, Kabatereine NB, Tchuem Tchuente LA, Rahbek C, Kristensen TK, 2012. Large-scale determinants of intestinal schistosomiasis and intermediate host snail distribution across Africa: does climate matter? *Acta Trop* (in press; doi:10.1016/j.actatropica.2011.11.010).
- Stockwell DRB, Noble IR, 1992. Induction of sets of rules from animal distribution data: a robust and informative method of data analysis. *Math Comput Simul* 33, 385-390.
- Teles HMS, Vaz JF, 1987. Distribuição de *Biomphalaria glabrata* (Say, 1818) (Pulmonata, Planorbidae) no Estado de São Paulo, Brasil. *Rev Saude Publica* 21, 508-512.
- Teodoro TM, Janotti-Passos LK, Carvalhos OS, Caldeira R, 2010. Occurrence of *Biomphalaria cousini* (Mollusca: Gastropoda) in Brazil and its susceptibility to *Schistosoma mansoni* (Platyhelminths: Trematoda). *Mol Phylogenet Evol* 57, 144-151.
- Waltari E, Hijmans RJ, Peterson AT, Nyári ÁS, Perkins SL, Guralnick RP, 2007. Locating Pleistocene refugia: comparing phylogeographic and ecological niche model predictions. *PLoS One* 2, e563.
- Wiley EO, McNyset KM, Peterson AT, Robins CR, Stewart AM, 2003. Niche modeling and geographic range predictions in the marine environment using a machine-learning algorithm. *Oceanography* 16, 120-127.