

A NEW MUTATION OF *PHEX* GENE IN A PATIENT WITH HYPERPHOSPHATURIA AND HYPERCALCIURIA

S. Carbonell Sala, A. Gozzini, A. Falchetti, E. Croppi, F. Cioppi, L. Masi, A. Tanini, M.L. Brandi

Department of Internal Medicine, University of Florence, Florence, Italy

The most common form of familial hypophosphatemic rickets (FHR), a group of disorders with similar clinical and biochemical features [hypophosphatemia, hyperphosphaturia, normal levels of 1,25(OH)₂D₃ and PTH, skeletal deformities, short stature, osteomalacia, dental abscesses bone pain], is represented by the dominant X-linked hypophosphatemic rickets (XLH). Individuals with FHR phenotype and a negative familial history in 60-80% of cases are carriers of mutations in *PHEX* gene, on chromosome Xp22.2-p22.1. The mice phenotypical analogue of the human XLH is represented by Hyp strand, in which a 3' deletion of *Phex* removes its COOH-terminal domain. The clinical consequences of *PHEX* inactivating mutations indicate that its encoded product, an endopeptidase member M13Zn-metalloproteinases family expressed at the skeletal level by osteoblasts, osteocytes, and odontoblasts, is involved in phosphate regulation and mineral homeostasis. *PHEX* inactivating mutations widespread along the gene cause XLH: exons 3-4-11-12-14-15-17-20-22 represent the regions with the higher rate of mutation; such mutations could enable the accumulation of phosphaturic factors and/or mineralization inhibitors. A 26 years old male patient (height 176cm, weight 65 kg) referred to our Centre exhibiting a clear hyperphosphaturia (>2000 mg/24h), hypophosphatemia, hypercalciuria (>600 mg/24h), hypocalcemia (□ 8,1 mg/die), PTH circulating levels at the upper values of the normal range and normal values of 25(OH)D; other symptoms were: deep asthenia, muscle pain and spasms, abundant diuresis (□ 2,5 lt/die). After obtaining the signed informed consent we performed a blood sampling from which genomic DNA has been prepared to analysed *PHEX* gene. The 22 exons and the intron-exon boundaries of *PHEX* gene have been investigated by a PCR/Sequencing protocol (ABI-Prism 3100). It has been determined the presence of a hemizygous missense mutation of *PHEX* gene in codon 401 (CCT/CTT) causing a Pro/Leu substitution in the extracellular domain closely a cysteine residue highly conserved in exon 11. Nearly future functional studies will be helpful to characterize the molecular mechanisms underlying this mutation.